

PRODUCT MONOGRAPH

RIVASA 81 MG EC

Acetylsalicylic Acid Delayed Release Tablets

81 mg

USP

Platelet aggregation inhibitor

Laboratoire Riva Inc.
660 Industriel Blvd.
Blainville, Quebec, Canada.
J7C 3V4
www.labriva.com

Date of revision :
December 21, 2015

Control number : 190401

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION.....	3
SUMMARY PRODUCT INFORMATION	3
INDICATIONS AND CLINICAL USE	3
CONTRAINDICATIONS	3
WARNINGS AND PRECAUTIONS	4
ADVERSE REACTIONS.....	6
DRUG INTERACTIONS	6
DOSAGE AND ADMINISTRATION	8
OVERDOSAGE.....	8
ACTION AND CLINICAL PHARMACOLOGY	10
STORAGE AND STABILITY	11
DOSAGE FORMS, COMPOSITION AND PACKAGING	11
PART II: SCIENTIFIC INFORMATION	12
PHARMACEUTICAL INFORMATION.....	12
CLINICAL TRIALS	13
DETAILED PHARMACOLOGY	16
TOXICOLOGY	19
PART III: CONSUMER INFORMATION.....	27

RIVASA 81 MG EC

Acetylsalicylic Acid Delayed Release Tablets, USP

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	Clinically Relevant Nonmedicinal Ingredients
Oral	Tablet / 81 mg	Not applicable <i>For a complete listing see Dosage Forms, Composition and Packaging section.</i>

INDICATIONS AND CLINICAL USE

RIVASA 81 mg EC (acetylsalicylic acid, ASA) is indicated for the following uses, based on its platelet aggregation inhibitory properties:

- for reducing the risk of vascular mortality in patients with a suspected acute myocardial infarction;
- for reducing the risk of a **first** non-fatal myocardial infarction in individuals deemed to be at sufficient risk of such an event by their physician.
 - There is no evidence for a reduction in the risk of **first** fatal myocardial infarction.
 - Acetylsalicylic acid does not reduce the risk of either cardiovascular mortality or **first** strokes, fatal or non-fatal.
 - The decrease in the risk of **first** non-fatal myocardial infarction must be assessed against a much smaller but not insignificant increase in the risk of haemorrhagic stroke as well as gastrointestinal bleeding.
- for reducing the risk of morbidity and death in patients with unstable angina and in those with previous myocardial infarction;
- for reducing the risk of transient ischemic attacks (TIA) and for secondary prevention of atherothrombotic cerebral infarction.

CONTRAINDICATIONS

- Patients who are hypersensitive to acetylsalicylic acid (ASA), salicylates, non-steroidal anti-inflammatory drugs (NSAIDs), analgesics, antipyretics or other ingredients in the product or component of the container. For a complete listing, see Dosage Forms, Composition and Packaging section of the product monograph.
- Acute gastrointestinal ulcer
- History of gastrointestinal ulcers
- Hemorrhagic diathesis
- Active or Severe hepatic failure, renal failure, or congestive heart failure

- Patients with a history of asthma induced by the administration of salicylates or substances with a similar action, notably NSAIDs
- Combination with methotrexate at doses of 15mg/week or more (see “Drug Interactions”).
- Last trimester of pregnancy

WARNINGS AND PRECAUTIONS

General

Acetylsalicylic acid (ASA) is one of the most frequent causes of accidental poisonings in toddlers and infants. Tablets should be kept well out of the reach of children.

ASA should be administered cautiously to patients with:

- uncontrolled hypertension
- impaired hepatic, renal function or cardiovascular circulation (e.g. renal vascular disease, congestive heart failure, volume depletion, major surgery, sepsis or major haemorrhagic events)
- a history of bleeding tendencies, significant anemia and/or hypofibrinogenemia
- concomitant treatment with anticoagulants (see “Drug Interactions”)
- concomitant treatment with ibuprofen in patients taking low-dose ASA (see “Drug Interactions”)

Hypersensitivity

Acetylsalicylic acid may precipitate bronchospasm and induce asthma attacks or other hypersensitivity reactions. Risk factors are present bronchial asthma, hay fever, nasal polyps, or chronic respiratory disease. This applies also for patients showing allergic reactions (e.g. cutaneous reactions, itching, urticaria) to other substances.

Hematologic

Due to effect on platelet aggregation, acetylsalicylic acid may be associated with an increased risk of bleeding. Caution is necessary when salicylates and anticoagulants are prescribed concurrently, as salicylates can depress the concentration of prothrombin in the plasma.

Peri-Operative Considerations

Due to its inhibitory effect on platelet aggregation which persists for several days after administration, acetylsalicylic acid may lead to an increased bleeding tendency during and after surgical operations (including minor surgeries, e.g. dental extractions).

Special Populations

Women attempting to conceive:

During the first and second trimester of pregnancy, acetylsalicylic acid containing drugs should not be given unless clearly necessary. If acetylsalicylic acid containing drugs are used by a woman attempting to conceive, or during the first and second trimester of pregnancy, the dose should be kept as low as possible and duration of treatment as short as possible.

Pregnant Women:

Acetylsalicylic acid inhibits prostaglandin synthesis. Inhibition of prostaglandin synthesis may adversely affect the pregnancy and/or the embryo/foetal development. Data from epidemiological

studies raise concern about an increased risk of miscarriage and of malformations after the use of a prostaglandin synthesis inhibitor in early pregnancy. The risk is believed to increase with dose and duration of therapy. Available data do not support any association between intake of acetylsalicylic acid and an increased risk for miscarriage. For acetylsalicylic acid the available epidemiological data regarding malformation are not consistent, but an increased risk of gastroschisis could not be excluded. A prospective study with exposure in early pregnancy (1st-4th month) of about 14,800 mother-child pairs has not yielded any association with an elevated rate of malformations.

During the third trimester of pregnancy, all prostaglandin synthesis inhibitors may expose the foetus to:

- cardiopulmonary toxicity (with premature closure of the ductus arteriosus and pulmonary hypertension);
- renal dysfunction, which may progress to renal failure with oligo-hydroamniosis;

Use of any prostaglandin synthesis inhibitors at the end of pregnancy may expose the mother and the child to:

- possible prolongation of bleeding time, an anti-aggregating effect which may occur even after very low doses
- inhibition of uterine contractions resulting in delayed or prolonged labour

Consequently, acetylsalicylic acid is contraindicated in the third trimester of pregnancy.

Nursing Women:

Acetylsalicylic acid and its metabolites pass into breast milk in small quantities. Since no adverse effects on the infant have been observed after occasional use, interruption of breast-feeding is usually unnecessary. However, on regular use or on intake of high doses, breast feeding should be discontinued early.

Pediatrics

A possible association between Reye's syndrome and the use of salicylates has been suggested but not established. Reye's syndrome has also occurred in many patients not exposed to salicylates. Acetylsalicylic acid should not be used in children and teenagers for viral infections with or without fever without consulting a physician. In certain viral illnesses, especially influenza A, influenza B and varicella, there is a risk of Reye's syndrome, a very rare but possibly life-threatening illness requiring immediate medical action. The risk may be increased when acetylsalicylic acid is given concomitantly; however, no causal relationship has been proven. Should persistent vomiting occur with such diseases; this may be a sign of Reye's syndrome.

Low Uric Acid Excretion:

At low doses, acetylsalicylic acid reduces excretion of uric acid. This can trigger gout in patients who already tend to have low uric acid excretion.

Glucose-6-phosphate dehydrogenase (G6PD) Deficiency:

In patient suffering from glucose-6-phosphate dehydrogenase (G6PD) deficiency, ASA may induce hemolysis or haemolytic anemia. Factors that may increase the risk of haemolysis are high dosage, fever, or acute infections.

Elderly

In general, ASA should be used with caution in elderly patients (≥ 60 years of age), as these patients may be more susceptible to adverse reactions.

Monitoring and Laboratory Tests

Salicylates can produce changes in thyroid function tests.

Isolated cases of liver function disturbances (transaminases increase) have been described.

ADVERSE REACTIONS

Many adverse reactions due to acetylsalicylic acid ingestion are dose-related. The following is a list of adverse reactions that have been reported in the literature and from both clinical and post-marketing experience.

Gastrointestinal (the frequency and severity of these adverse effects are dose-related): nausea, vomiting, diarrhea, gastrointestinal bleeding and/or ulceration, dyspepsia, heartburn, hematemesis, melena, abdominal pain, and rarely gastrointestinal inflammation.

Bleeding: Due to platelet inhibition, bleedings e.g. perioperative haemorrhage, hematomas, epistaxis, urogenital bleedings, and gingival bleedings may occur.

Serious bleedings, such as gastrointestinal tract hemorrhages, and cerebral hemorrhages are rare. Isolated cases of potentially life threatening bleedings have been reported, especially in patients with uncontrolled hypertension and/or concomitant antihemostatic agents.

Ear: dizziness, tinnitus, vertigo, hearing loss. Dizziness and tinnitus have been reported, which may be indicative of an overdose.

Hematologic: leukopenia, thrombocytopenia, purpura, anemia. Anemia with respective laboratory and clinical signs and symptoms, such as asthenia, pallor, and hypoperfusion is generally caused by bleeding (e.g. occult microbleeding, acute or chronic bleeding). Hemolysis and hemolytic anemia in patients with severe forms of glucose-6-phosphate dehydrogenase (G6PD) deficiency has been reported.

Dermatologic and hypersensitivity: urticaria, pruritus, skin eruptions, asthma, anaphylaxis, edema, nasal congestion and rhinitis. Severe allergic reactions, including anaphylactic shock are very rarely reported.

Miscellaneous: mental confusion, drowsiness, sweating, thirst. Transient hepatic impairment with increase in liver transaminases has very rarely been reported. Renal impairment and acute renal failure have been reported.

DRUG INTERACTIONS

Overview

Acetylsalicylic acid should be used with caution with other products that have anticoagulation or antiplatelet effects, as these effects may be potentiated. Drugs that bind to protein binding sites should also be used cautiously since acetylsalicylic acid may displace drugs from their protein binding site.

Contraindicated Interactions

Methotrexate, used at doses of 15 mg/week or more: Increased hematological toxicity of methotrexate (due to decreased renal clearance of methotrexate by anti-inflammatory agents in general and displacement of methotrexate from its plasma protein binding by salicylates). See

“Contraindications”.

Drug-Drug Interactions

Methotrexate, used at 15mg/week or less: Salicylates may retard the elimination of methotrexate by decreasing renal clearance of methotrexate, displacing methotrexate from protein binding sites, and thereby increasing its hematological toxicity.

Anti-coagulants, thrombolytics / other inhibitors of platelet aggregation / hemostasis, e.g. warfarin, heparin: Caution is necessary when salicylates and anticoagulants, thrombolytics / other inhibitors of platelet aggregation / hemostasis prescribed concurrently, as salicylates can depress the concentration of prothrombin in the plasma, leading to an increased risk of bleeding.

Oral hypoglycemics, e.g. insulin, sulfonylureas: Large doses of salicylates have a hypoglycemic action and may enhance the effect of oral hypoglycemic agents. Diabetics receiving concurrent salicylate and hypoglycemic therapy should be monitored closely: reduction of the sulfonylurea hypoglycemic drug dosage may be necessary.

Diuretics: Diuretics in combination with acetylsalicylic acid at higher doses leads to decreased glomerular filtration via decreased prostaglandin synthesis. As a result, sodium excretion may be decreased by salicylate administration.

Uricosuric Agents: Salicylates in large doses are uricosuric agents; smaller amounts may depress uric acid clearance and thus decrease the uricosuric effects of other drugs.

Valproic Acid: Salicylates may alter valproic acid (VPA) metabolism and may displace VPA from protein binding sites, possibly intensifying the effects of VPA. Caution is recommended when VPA is administered concomitantly with salicylates.

Glucocorticoids (systemic), except hydrocortisone used as replacement therapy in Addison’s disease: Decreased blood salicylate levels during corticosteroid treatment and risk of salicylate overdose after this treatment is stopped via increased elimination of salicylates by corticosteroids.

Angiotensin Converting Enzyme (ACE) Inhibitors: The hyponatremic and hypotensive effects of ACE inhibitors *may* be diminished by the concomitant administration of acetylsalicylic acid due to its indirect effect on the renin-angiotensin conversion pathway (i.e. inhibition of vasodilatory prostaglandins leading to decreased glomerular filtration). The potential interaction may be related to the dose of acetylsalicylic acid (3g/day or more).

Selective Serotonin Re-uptake Inhibitors (SSRIs): Increased risk of upper gastrointestinal bleeding due to possibly synergistic effect.

Digoxin: Plasma concentrations of digoxin are increased due to a decrease in renal excretion.

ASA and other NSAIDs: The use of other non-steroidal anti-inflammatory drugs (NSAIDs) with salicylates at high doses ($\geq 3\text{g/day}$) may increase the risk of ulcers and gastrointestinal bleeding due to a synergistic effect.

Ibuprofen: Ibuprofen can interfere with the anti-platelet effect of low dose acetylsalicylic acid (81-325

mg per day). Long-term daily use of ibuprofen may render ASA less effective when used for cardioprotection and stroke prevention. To minimize this interaction, regular users of ibuprofen and of low-dose, immediate-release ASA should take the ibuprofen at least one hour after and 11 hours before the daily ASA dose. The use of delayed-release (e.g. enteric-coated) ASA is not recommended when using ibuprofen regularly. Healthcare professionals should advise consumers and patients regarding the appropriate concomitant use of ibuprofen and ASA.

Drug-Food Interactions

Interactions with food have not been established.

Drug-Herb Interactions

Interactions with herb have not been established.

Drug-Laboratory Interactions

Salicylates can produce changes in thyroid function tests.

Drug-Lifestyle Interactions

Alcohol: Increased damage to gastrointestinal mucosa and prolonged bleeding time due to additive effects of acetylsalicylic acid and alcohol. Patients having 3 or more alcoholic drinks per day should consult their physician before use.

DOSAGE AND ADMINISTRATION

RIVASA 81 mg EC tablets should preferably be taken after meals, with plenty of liquid.

Dosing Considerations

Please see below for specific dosing instructions for each indication.

Recommended Dose and Dosage Adjustment

Platelet aggregation inhibitor:

Suspected Acute Myocardial Infarction: An initial dose of at least 162 mg chewed or crushed to ensure rapid absorption as soon as a myocardial infarction is suspected. The same dose should be given as maintenance over the next 30 days. After 30 days, consider further therapy based on dosage and administration for prevention of recurrent MI (see Prior Myocardial Infarction).

Prevention of a first non-fatal myocardial infarction: 81 - 325 mg once daily, according to the individual needs of the patient, as determined by the physician.

Prior Myocardial Infarction or Unstable Angina Pectoris: 81 - 325 mg daily according to the individual needs of the patient, as determined by the physician.

Transient Ischemic Attack and Secondary Prevention of Atherothrombotic Cerebral Infarction: 81 - 325 mg daily according to the individual needs of the patient, as determined by the physician.

OVERDOSAGE

Mild Overdose or Early Poisoning - burning in the mouth, lethargy, nausea, vomiting, tinnitus, sweating, thirst, tachycardia or dizziness.

Moderate Overdose - all of the symptoms from mild overdose plus tachypnea, hyperpyrexia, sweating, dehydration, loss of coordination, restlessness, mental confusion.

Severe Overdose - all of the symptoms from moderate overdose plus hypotension, hallucinations, stupor, hypoglycemia, convulsions, cerebral edema, oliguria, renal failure, cardiovascular failure, coma, hemorrhage, metabolic acidosis, respiratory alkalosis and/or failure.

Emergency Management:

- 1- Immediate transfer to hospital and maintain cardiovascular and respiratory support.
- 2- Gastric lavage, administration of activated charcoal,
- 3- Check of acid-base balance and correct if necessary.
- 4- Alkaline diuresis so as to obtain urine pH between 7.5 and 8 should be considered when plasma salicylate concentration is greater than 500mg/L (3.6 mmol/L) in adults or 300mg/L (2.2 mmol/L) in children
- 5- Hemodialysis should be considered in severe poisoning 800mg/L (5.8 mmol/L) in adults and 700mg/L (5.0 mmol/L) in children, as renal elimination of salicylates may be slow due to the presence of acidic urine and renal failure. Hemodialysis should also be considered if the patient is experiencing severe systemic metabolic acidosis (arterial pH < 7.2), acute renal failure, pulmonary edema or CNS symptoms such as: drowsiness, agitation, coma or convulsions.
- 6- Fluid losses should be replaced with hypotonic solution (e.g. half saline) and supplemented with glucose 50 to 100g/L.
- 7- Symptomatic treatment.

Fatal Dose: varies from 10 to 30g of acetylsalicylic acid. However, (in one case) 130g of acetylsalicylic acid was ingested without fatal outcome.

For the management of a suspected drug overdose, contact your regional Poison Control Centre.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

Acetylsalicylic acid interferes with the production of prostaglandins in various organs and tissues through acetylation of the enzyme cyclo-oxygenase. Prostaglandins are themselves powerful irritants and produce headaches and pain on injection in man. Prostaglandins also appear to sensitize pain receptors to other noxious substances such as histamine and bradykinin. By preventing the synthesis and release of prostaglandins in inflammation, acetylsalicylic acid may avert the sensitization of pain receptors.

The antipyretic activity of acetylsalicylic acid is due to its ability to interfere with the production of prostaglandin E₁ in the brain. Prostaglandin E₁ is one of the most powerful pyretic agents known.

The inhibition of platelet aggregation by acetylsalicylic acid is due to its ability to interfere with the production of thromboxane A₂ within the platelet. Thromboxane A₂ is, largely, responsible for the aggregating properties of platelets.

In vitro studies have shown that acetylsalicylic acid enhances the activity of the Nitric oxide (NO)-cGMP system and heme oxygenase-1 (HO-1) by acting on endothelial NO synthase site.

Pharmacokinetics

Absorption:

When acetylsalicylic acid is taken orally, it is rapidly absorbed from the stomach and proximal small intestine. The gastric mucosa is permeable to the non-ionized form of acetylsalicylic acid, which passes through the stomach wall by a passive diffusion process.

Optimum absorption of salicylate in the human stomach occurs in the pH range of 2.15 to 4.10. Absorption in the small intestine occurs at a significantly faster rate than in the stomach. After an oral dose of 0.65 g acetylsalicylic acid, the plasma acetylsalicylate concentration in man usually reaches a level between 0.6 and 1.0 mg % in 20 minutes after ingestion and drops to 0.2 mg % within an hour. Within the same period of time, half or more of the ingested dose is hydrolyzed to salicylic acid by esterases in the gastrointestinal mucosa and the liver, the total plasma salicylate concentration reaching a peak between one or two hours after ingestion, averaging between 3 and 7 mg %. Many factors influence the speed of absorption of acetylsalicylic acid in a particular individual at a given time; tablet disintegration, solubility, particle size, gastric emptying time, psychological state, physical condition, nature and quantity of gastric contents, etc., all affect absorption.

Distribution:

Distribution of salicylate throughout most body fluids and tissues proceeds at a rapid rate after absorption. Aside from the plasma itself, fluids which have been found to contain substantial amounts of salicylate after oral ingestion include spinal, peritoneal and synovial fluids, saliva and milk. Tissues containing high concentrations of the drug are the kidney, liver, heart and lungs. Concentrations in the brain are usually low, and are minimal in feces, bile and sweat.

The drug readily crosses the placental barrier. At clinical concentrations, from 50% to 90% of the salicylate is bound to plasma proteins especially albumin, while acetylsalicylic acid itself is bound to only a very limited extent. However, acetylsalicylic acid has the capacity of acetylating various proteins,

hormones, DNA, platelets and hemoglobin, which at least partly explains its wide-ranging pharmacological actions.

Metabolism:

The liver appears to be the principal site for salicylate metabolism, although other tissues may also be involved. The three chief metabolic products of acetylsalicylic acid or salicylic acid are salicyluric acid, the ether or phenolic glucuronide and the ester or acyl glucuronide. A small fraction is also converted to gentisic acid and other hydroxybenzoic acids. The half-life of acetylsalicylic acid in the circulation is from 13 to 19 minutes so that the blood level drops quickly after absorption is complete. However, the half-life of the salicylate ranges between 3.5 and 4.5 hours, which means that 50% of the ingested dose leaves the circulation within that time.

Excretion:

Excretion of salicylates occurs principally via the kidney, through a combination of glomerular filtration and tubular excretion, in the form of free salicylic acid, salicyluric acid, as well as phenolic and acyl glucuronides. Salicylate can be detected in the urine shortly after its ingestion but the full dose requires up to 48 hours for complete elimination. The rate of excretion of free salicylate is extremely variable, reported recovery rates in human urine ranging from 10% to 85%, depending largely on urinary pH. In general, it can be stated that acid urine facilitates reabsorption of salicylate by renal tubules, while alkaline urine promotes excretion of the drug.

With the administration of 325mg, elimination of acetylsalicylic acid is linear following a first order kinetics. At high concentrations, elimination half life increases.

Special Populations and Conditions:

Absorption and clearance of salicylates are not affected by gender or age.

STORAGE AND STABILITY

Store between 15-30°C.

DOSAGE FORMS, COMPOSITION AND PACKAGING

RIVASA 81 mg EC: Each round, convex tablet is blue in colour debossed with a "V" on one side and plain on the other side. The tablet composition includes 81 mg of acetylsalicylic acid and the non-medicinal ingredients: Colloidal silica, corn starch, FD&C blue #1, hypromellose, methacrylic acid copolymer, microcrystalline cellulose, purified water, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate.

RIVASA 81 mg EC is packaged in bottles with child resistant caps and counts of 1000.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION

Drug Substance


Proper name: acetylsalicylic acid

Chemical name: 2-(Acetyloxy) benzoic acid; salicylic acid acetate

Molecular formula: C₉H₈O₄

Molecular Mass: 180.16 g/mol

Structural formula:



Physicochemical properties:

Description: White granules, commonly tabular or needle-like, or white crystalline powder. Odourless or having a faint odour.

Solubility: Slightly soluble in water; freely soluble in alcohol; soluble in chloroform and ether; sparingly soluble in absolute ether.

pK value (25°C): 3.49

Melting Point: 135° C (rapid heating)

CLINICAL TRIALS

Comparative Bioavailability Studies

A single-dose, randomized, double-blinded, crossover, pivotal comparative bioavailability study was conducted comparing Enteric-Coated Daily Low Dose RIVASA 81 mg EC Tablets (acetylsalicylic acid delayed release tablets) (Laboratoire Riva Inc.) and Aspirin® 81 mg Tablets (acetylsalicylic acid delayed release tablets) (Bayer Inc., Consumer Care) in 39 healthy male and female volunteers under fasting conditions.

Salicylic Acid (2 x 81 mg) From measured data Geometric Mean Arithmetic Mean (CV %)				
Parameter	^a Test	^b Reference	% Ratio of Geometric Means	90 % Confidence Interval
AUC _T (ng.h/mL)	40035.803 41180.528 (24.94)	40009.476 40905.966 (22.11)	100.07	96.84 - 103.39
AUC _I (ng.h/mL)	41086.748 42459.504 (25.62)	40672.101 41148.135 (23.10)	101.02	97.56 - 104.60
C _{max} (ng/mL)	8345.017 8704.618 (28.85)	8824.099 8999.336 (19.75)	94.57	85.90 - 104.12
T _{max} [§] (h)	4.50 (1.57 - 10.00)	4.50 (2.50 - 7.00)		
T _½ ^ε (h)	2.39 (30.86)	2.47 (52.17)		

^aEnteric Coated Daily Low Dose RIVASA 81 mg EC (acetylsalicylic acid delayed release tablets) (Laboratoire Riva Inc.)

^bAspirin® 81 mg (acetylsalicylic acid delayed release tablets) (Bayer Inc., Consumer Care) were purchased in Canada.

[§]Expressed as median (range) only.

^εExpressed as the arithmetic mean (CV%) only

Comparative Bioavailability Studies

A single-dose, randomized, double-blinded, crossover, pivotal comparative bioavailability study was conducted comparing Enteric-Coated Daily Low Dose RIVASA 81 mg EC Tablets (acetylsalicylic acid delayed release tablets) (Laboratoire Riva Inc.) and Aspirin® 81 mg Tablets (acetylsalicylic acid delayed release tablets) (Bayer Inc., Consumer Care) in 37 healthy male and female volunteers under fed conditions.

Salicylic Acid (2 x 81 mg) From measured data Geometric Mean Arithmetic Mean (CV %)				
Parameter	^a Test	^b Reference	%Ratio of Geometric Means	90 % Confidence Interval
AUC _T (ng.h/mL)	41174.624 44213.823 (38.45)	44138.779 46949.222 (37.27)	93.28	88.13 - 98.74
AUC _I (ng.h/mL)	41175.322 43247.467 (26.55)	42513.535 44344.611 (36.63)	96.85	91.18 - 102.88
C _{max} (ng/mL)	6555.507 7025.088 (32.21)	7475.813 7875.047 (32.88)	87.69	79.60 - 96.60
T _{max} [§] (h)	9.00 (3.00 - 24.05)	10.00 (4.60 - 24.00)		
T _½ ^ε (h)	2.52 (19.18)	2.74 (35.63)		

^aEnteric Coated Daily Low Dose RIVASA 81 mg EC (acetylsalicylic acid delayed release tablets) (Laboratoire Riva Inc.)

^bAspirin® 81 mg (acetylsalicylic acid delayed release tablets) (Bayer Inc., Consumer Care) were purchased in Canada.

[§]Expressed as median (range) only.

^εExpressed as the arithmetic mean (CV%) only

Study demographics and trial design

Anti-Platelet Aggregation Studies

Study #/ cross- reference	Trial Design	Dosage, route of administration and duration	Study Subjects (n = number)	Mean age (Range)	Gender
Indication: Reducing the risk of vascular mortality in patients with a suspected acute myocardial infarction.					
ISIS – 2 Ref 102	Multicentre international 2x2 factorial, randomized placebo controlled study.	160-162.5 mg oral for 30 days after suspected acute MI. (Median follow-up to 15 months).	ASA 8587, Streptokinase 8592, ASA + Strep 4292, Placebo 4300	Not available	Not available
Indication: Reducing the risk of a first non-fatal myocardial infarction in individuals deemed to be at sufficient risk of such an event by their physician					
TPT Ref 61	Randomized, factorial, placebo-controlled, parallel-group study	warfarin (mean) 4.1mg, ASA 75mg	warfarin + ASA 1,277 warfarin + ASA placebo 1,268 ASA + warfarin placebo 1,268 ASA placebo + warfarin placebo 1,272	45-69 years	Male
HOT Ref 101	Prospective, randomized, open with blinded endpoint evaluation (PROBE). ASA component was double blinded	ASA 75mg or placebo; felodine 5mg, ACE-inhibitors, β Z blockers, diuretics mean - 3.8 years	19,567 subjects of which 18,790 were randomized to ASA or Placebo (ASA = 9,399; Placebo = 9,391)	61.5 years - mean (50-80 years)	Male 53% Female 47%
PHS Ref 100	Double blind placebo controlled, 2x2 factorial randomized parallel group	ASA 325mg every other day for 60.2 months	22,071 ASA = 11,037 Placebo = 11,034	40 to 84 years	Male
Indication: Reducing the risk of morbidity and death in patients with unstable angina and in those with previous myocardial infarction					
RISC Ref 76	Prospective randomized, double blind, placebo controlled, multicentre study	ASA 75mg daily for 3 months after initial heparin by IV for 5 days	-Heparin 198 -ASA 189 -Heparin + ASA 210 - Placebo 199	58 years	Male
RISC Trial, 12 month follow-up Ref 89	Prospective randomized, double blind, placebo controlled, multicentre study	ASA 75mg daily for 3 months after initial IV heparin for 5 days	-Heparin 198 -ASA 189 -Heparin + ASA 210 - Placebo 199	58 years	Male
Verheugt et al. Ref 88	Prospective, randomized, placebo-controlled, comparative multicentre study	ASA 100mg for approx. 3 months	ASA 50 Placebo 50	ASA 61 years Placebo 64 years	ASA 72% male Placebo 76% male
SAPAT Ref 54	Prospective, randomized, double blind placebo controlled, multicentred study	ASA 75mg daily for up to 6 years (median 50 months)	ASA 1009 Placebo 1026	52 years	ASA male 51% Placebo males 53%
Indication: Reducing the risk of transient ischemic attacks (TIA) and for secondary prevention of atherothrombotic cerebral infarction					

Study #/ cross- reference	Trial Design	Dosage, route of administration and duration	Study Subjects (n = number)	Mean age (Range)	Gender
SALT Ref 81	Prospective, randomized, double blind, placebo controlled, multicentre study	ASA 75mg daily for minimum of 12 months and maximum of 63 months (mean 30.6 months)	ASA 676 Placebo 684	50-79 years ASA mean: 67 years PLA mean: 66.8 years	ASA 65.4% male Placebo 66.2% male
Lindblad et al. Ref 57	Prospective, randomized, double blind placebo controlled study	ASA 75mg daily for 6 months	ASA 117 Placebo 115	66 years (40-81 years)	75% male

Study results

Platelet Aggregation Studies Results

Study #	Primary Endpoints	Associated value and statistical significance for ASA compared to Placebo		
Indication: Reducing the risk of vascular mortality in patients with a suspected acute myocardial infarction.				
		Value	ASA vs. Placebo	
ISIS – 2 Ref 102	Vascular death after 5 week period	ASA 9.4%, Placebo 11.8% Odds reduction 23%	2p < 0.00001 ASA was statistically significantly better than placebo	
Indication: Reducing the risk of a first non-fatal myocardial infarction in individuals deemed to be at sufficient risk of such an event by their physician				
TPT Ref 61	All ischemic heart disease defined as the sum of fatal and non-fatal events (i.e. coronary death and fatal and non-fatal myocardial infarction).	ASA 10.2%, Placebo 13.3% 20% reduction in IHD	p = 0.04 ASA was statistically significantly better than placebo	
HOT Ref 101	Major cardiovascular events were defined as all (fatal and non-fatal) myocardial infarctions, all (fatal and non-fatal) strokes, and all other cardiovascular deaths.	Reduction in all cardiovascular events by 15 % and all myocardial infarction by 36%	p=0.03 p = 0.002 ASA was statistically significantly better than placebo	
PHS Ref 100	fatal and non-fatal myocardial infarction	325 mg ASA every other day: 44% reduction in risk of MI in ASA vs. Placebo group Relative Risk 0.56, 95% CI 0.45-0.70	p<0.00001 P<0.0001 ASA was statistically significantly better than placebo	
Indication: Reducing the risk of morbidity and death in patients with unstable angina and in those with previous myocardial infarction				
		Value	ASA vs. Placebo	ASA vs. Comparator

Study #	Primary Endpoints	Associated value and statistical significance for ASA compared to Placebo		
RISC	Death or non-fatal MI	5 days: Risk Ratio 0.43 (CI 0.21-0.91) 30 days: Risk Ratio 0.31 (CI 0.18-0.53) 90 days: Risk Ratio 0.36 (0.21-0.57)	p=0.03 p<0.0001 p<0.0001 ASA was statistically significantly better than placebo	Heparin was not statistically significantly better than placebo and there was no comparison to ASA
RISC Trial, 12 month follow-up	MI and death	6 months: ASA-35 events, heparin 76 events. Risk Ratio 0.46 (CI 0.31-0.67) 12 months, ASA 44 events, heparin 85 events. Risk Ratio 0.52 (CI 0.37-0.72)	p<0.0001 p=0.0001 ASA was statistically significantly better than placebo	Not Performed
Verheugt et al.	Reinfarction rate	ASA 2 patients (4%), Placebo 9 patients (18%)	p<0.03 ASA was statistically significantly better than placebo	Not Performed
SAPAT	non-fatal or fatal MI or sudden death	ASA 8%, Placebo 12%	p=0.003 ASA was statistically significantly better than placebo	Not Performed
Indication: Reducing the risk of transient ischemic attacks (TIA) and for secondary prevention of atherothrombotic cerebral infarction				
		Value	ASA vs. Placebo	
SALT	Risk of stroke or death	18 % reduction in risk: Relative Risk 0.82 (CI 0.67-0.99)	p=0.02 ASA was statistically significantly better than placebo	
Lindblad et al.	Stroke (without complete recovery) at 6 months	ASA 2 cases, Placebo 11 cases	p=0.01 ASA was statistically significantly better than placebo	

DETAILED PHARMACOLOGY

Effects on platelets: relation to hemostasis and thrombosis.

Platelets play an important role in normal hemostasis and clinical pathologic and experimental evidence indicates that their aggregation may play an equally important role in the evolution of a variety of disease states including cerebrovascular disease, ischemic heart disease and myocardial infarction. Acetylsalicylic acid inhibits platelet aggregation by irreversibly acetylating platelet cyclooxygenase, thereby blocking the production of prostaglandin endoperoxides PGG₂ and PGH₂ which are precursors of the major platelet-aggregating material, thromboxane A₂, which is also a powerful vasoconstrictor. However, acetylsalicylic acid does not prevent the adherence of platelets to damaged vessel walls or the release of granule contents from these adherent platelets. As the anuclear platelets are unable to synthesize new enzyme molecules to replace those that have been inactivated, inhibition of platelet

aggregation by acetylsalicylic acid thus persists for the life of the platelets. Daily administration of 20 to 40 mg of acetylsalicylic acid to healthy volunteers reduced platelet thromboxane production but inhibited platelet aggregation only partially. When administered to patients recovering from myocardial infarction, 50 mg acetylsalicylic acid daily had the same effects on thromboxane production, platelet aggregation and bleeding times as 324 mg daily. Other studies show that acetylsalicylic acid doses of 40 to 325 mg daily suppressed thromboxane production by at least 80%, but 80 mg acetylsalicylic acid daily was the lowest dose required for maximum cumulative thrombocyte function inhibition. The protective effect of acetylsalicylic acid against experimentally induced thrombosis or atherosclerosis has been demonstrated in several animal models.

Besides inhibiting the biosynthesis of thromboxane A₂ by platelets, acetylsalicylic acid also interferes with the production of prostacyclin (PGI₂) by vascular endothelial cells, the above-mentioned prostaglandin endoperoxides being common precursors of both thromboxane A₂ and prostacyclin. This latter compound is one of the most powerfully acting platelet deaggregators and vasodilators and thus it would appear that the interference with the hemostatic processes by acetylsalicylic acid depends on the thromboxane-prostacyclin balance. In fact, it has been suggested that under some conditions, high doses of acetylsalicylic acid may be thrombogenic. However, in contrast to platelets, the vascular endothelial cells are able to regenerate cyclo-oxygenase in a relatively short time and therefore therapeutic doses of acetylsalicylic acid are likely to produce a lesser inhibition of the vascular prostacyclin system than of the platelet thromboxane-forming mechanism. In fact, there is no clinical evidence to indicate that high doses of acetylsalicylic acid would result in an increased risk of thromboembolism. Indeed, quite the contrary was observed and, in a controlled study, paradoxical shortening of the bleeding time was not observed at a daily acetylsalicylic acid dose of 3.6 g. Lower dosages of acetylsalicylic acid make selective blocking of the TxA₂-synthesis without a simultaneous blocking of PGI₂-production possible.

The use of acetylsalicylic acid in patients with a suspected acute myocardial infarction was investigated in a large multi-centre trial involving over 17,000 patients. Treatment with acetylsalicylic acid resulted in a 23% reduction in the risk of vascular mortality versus placebo at 5 weeks. This use translates to a reduction of 24 deaths and 14 non-vascular events per 1000 patients treated. The effect of time to therapy revealed that patients treated with acetylsalicylic acid “early” (0 to 4 hours) versus “late” (5 to 24 hours) after symptom onset experienced reductions in the odds of vascular death of 25% versus 21%, versus placebo at 5 weeks. ‘Early’ treatment with acetylsalicylic acid resulted in the saving of 4 additional lives per 1000 patients versus ‘late’ treatment.

Long term follow-up (up to 10 years) of patients in this study established that the early survival advantage to acetylsalicylic acid persisted long term, and that this prolonged benefit was additive to that of fibrinolytic therapy.

The use of acetylsalicylic acid for secondary prevention of thrombotic events is supported by a comprehensive overview of a number of clinical trials involving patients who already had some type of vascular disease (myocardial infarction, unstable angina, stroke or transient cerebral ischemia). Overall, these studies point to a 26-28 % reduction of the combined endpoints of MI, stroke, or vascular deaths by treatment with acetylsalicylic acid alone at doses of 75 to 325 mg daily. Studies which directly compared low doses with higher doses (30-1200 mg/day), indicated that the incidence of gastrointestinal adverse effects were significantly less common with the lower doses.

In a study in patients undergoing coronary artery bypass surgery (CABG), patients given acetylsalicylic acid at a dosage of 80 mg to 650 mg within 48 hours of revascularization had a risk of dying reduced to 1.3% as compared to 4.0% for those who did not receive treatment (P<0.001). There was a reduction in the incidence of myocardial infarction of 2.8% vs. 5.4%, p < 0.001. In total, the reduction in fatal and

non-fatal outcomes was lower in those who received ASA, 10.6% vs. 18.6% in those who did not ($p < 0.001$). The investigators Perioperative Ischemia Research Group (PIRG) concluded that early use of acetylsalicylic acid after coronary by-pass surgery is safe and is associated with a reduce risk of death and ischemic complications involving the heart, brain, kidneys and gastrointestinal tract. There was no acetylsalicylic acid dose effect observed for either fatal or non-fatal outcomes with total doses lower than 325mg daily.

Recent discussions have focused on the efficacy of acetylsalicylic acid for the primary prevention of myocardial infarction and stroke. Two large scale randomized trials, aimed at evaluating prophylactic use of acetylsalicylic acid, were conducted among apparently healthy male physicians (22,000 in the United States and 5,000 in the United Kingdom) and their results have been published. In the summary overview of the combined results presented by the principal investigators, the authors state that: "Taken together, these two primary prevention studies demonstrate a significant ($p < 0.0001$) reduction in non-fatal myocardial infarction of about one third."

On the other hand, the same two studies have not indicated any reduction in overall vascular mortality and also suggested a slight increase in the risk of non-fatal disabling stroke. Current controversy exists about the applicability of these findings, obtained in a selected population, to the general public. As well, the optimum dosage regimen still remains an open question in this regard. Thus, the use of acetylsalicylic acid for primary prevention should remain, in the words of the principal investigators: "a matter of judgment in which the physician considers the cardiovascular risk profile of the patient and balances the known hazards of ASA...against the clearly established reduction in the incidence of a first myocardial infarction".

Effect of Ibuprofen on Platelet Aggregation, Bleeding and Clotting Times in Normal Volunteers.

Experimental data suggest that ibuprofen may inhibit the effect of low dose ASA (81-325 mg per day) on platelet aggregation when they are dosed concomitantly. In one study, when a single dose of ibuprofen 400mg was taken within 8 hours before or within 30 minutes after immediate release ASA dosing (81mg), a decreased effect of acetylsalicylic acid on the formation of thromboxane or platelet aggregation occurred. However, the limitations of these data and the uncertainties regarding extrapolation of *ex vivo* data to the clinical situation imply that no firm conclusions can be made for regular ibuprofen use, and no clinically relevant effect is considered to be likely for occasional ibuprofen use. In a more recent double blind, randomized, placebo-controlled trial with healthy subjects by Cryer et. al, 2005, it has been shown that the drug-drug interaction is absent when immediate release ASA (81 mg) was taken 1 hour before taking ibuprofen (400 mg, TID) and also when ibuprofen was given 11 hours before the intake of low dose ASA. Thus, in order to adequately minimize potential interaction, the recommended dosing schedule for immediate release low dose ASA is to wait at least 11 hours after or 1 hour before taking up to a 400mg dose of ibuprofen.

MICROBIOLOGY

Not applicable

TOXICOLOGY

The clinical and pathological signs of poisoning from toxic and lethal oral doses of acetylsalicylic acid have been extensively described for man, much less extensively for other species.

The acute toxicity of acetylsalicylic acid in animals has been studied and reviewed in detail by Boyd. The signs of poisoning in rats from doses in the lethal range are due to varying degrees of gastroenteritis, hepatitis, nephritis, pulmonary edema, encephalopathy, shock and minor toxic effects on other organs and tissues. Death is due to convulsions or cardiovascular shock. The major difference between species appears to be the ability to vomit toxic doses seen in man, cats and dogs, but not in mice, rats and rabbits. Otherwise, the pathological reaction to toxic doses of acetylsalicylic acid is similar in all species in which such studies have been reported. The acute oral LD₅₀ values have been reported as being over 1.0 g/kg in man, cat and dog, 0.92 g/kg in female and 1.48 g/kg in male albino rats, 1.19 g/kg in guinea pig, 1.1 g/kg in mouse and 1.8 g/kg in rabbit.

Chronic toxicity studies were reported in mice and rats. When acetylsalicylic acid was administered at 2 to 20 times the maximum tolerated clinical dose to mice for up to one year, a dose-related deleterious effect was observed on mean survival time, number of young born and number of young raised to weaning age. No evidence of carcinogenic effect was found.

The chronic oral LD₅₀ in male albino rats has been reported as 0.24 g/kg/day when given for 100 days. At these daily doses acetylsalicylic acid produced no anorexia and no loss of body weight. It did produce polydipsia, aciduria, diuresis, drowsiness, hyperreflexia, piloerection, rapid and deep respiration, tachycardia, and during the second month, soft stools, epistaxis, sialorrhea, dacryorrhea and death in hypothermic coma. Autopsy disclosed the presence of a hypertrophied stomach, renal congestion, mild hepatitis and pneumonitis. While teratogenic effects were noted in animals at near lethal doses, there is no evidence to indicate that acetylsalicylic acid is teratogenic in man.

REFERENCES

- Abbott F, Kassam J, Orr J, and K. Farrell. The effects of Aspirin on valproic acid metabolism. *Clin. Pharmacol. Ther.* 1986;40:94-100.
- Ali NAJ, Al-Naama LM, Khalid LO. Haemolytic potential of three chemotherapeutic agents and aspirin in glucose-6-phosphate dehydrogenase deficiency. *East Mediterr Health J.* 1999 May;5(3):457-64.
- Altman R, Bouillon F, Rouvier J, Raca R, de la Fuente, Favaloro R. Aspirin and prophylaxis of thromboembolic complications in patients with substitute heart valves. *J Thorac Cardiovasc Surg* 1976; 72: 127-9.
- Amrein PC, Ellman L, Harris WH. Aspirin prolongation of bleeding time and perioperative blood loss. *JAMA* 1981; 245: 1825-8.
- Analgesic-antipyretic and anti-inflammatory agents: the salicylates. *In: Goodman and Gilman's, The Pharmacological Basis of Therapeutics*, 9th Edition, Hardman JG, Limbird LE, Gilman (eds), McGraw-Hill, USA, 1996.
- Antiplatelet Trialists' Collaboration. Collaborative overview of randomised trials of antiplatelet therapy - I: Prevention of death, myocardial infarction, and stroke by prolonged antiplatelet therapy in various categories of patients. *Br Med J* 1994; 308: 81-106.
- Antiplatelet Trialists' Collaboration. Collaborative overview of randomised trials of antiplatelet therapy - II: Maintenance of vascular graft or arterial patency by antiplatelet therapy. *Br Med J* 1994; 308: 158-68.
- Antiplatelet Trialists' Collaboration. Collaborative overview of randomised trials of antiplatelet therapy - III: Reduction in venous thrombosis and pulmonary embolism by antiplatelet prophylaxis among surgical and medical patients. *Br Med J* 1994; 308: 235-46.
- Antiplatelet Trialists' Collaboration. Secondary prevention of vascular disease by prolonged antiplatelet treatment. *Br Med J* 1988; 296: 320-31.
- Antonucci R, Zaffanello M, Puxeddu E, Porcella A, Cuzzolin L, Pilloni MD, Fanos V. Use of non-steroidal anti-inflammatory drugs in pregnancy: impact on the fetus and newborn. *Curr Drug Metab.* 2012 May 1;13(4):474-90.
- Aspirin Myocardial Infarction Study Research Group. A randomized controlled trial of Aspirin in persons recovered from myocardial infarction. *JAMA* 1980; 243: 661-9.
- Aspirin Myocardial Infarction Study Research Group: The Aspirin myocardial infarction study: final results. *Circulation* 1980; 62 (Suppl V): V79-V84.
- Bailey JM. Prostacyclins, thromboxane and cardiovascular disease. *Tr Biochem Sci* 1979; 4: 68-71.
- Baxter, K. *Stockley's Drug Interactions*, 9 Edition. Royal Pharmaceutical Society, 2010; 1123, 1135-1136
- Bayer Inc. Product Monograph. ASPIRIN® Regular Strength; ASPIRIN® Extra Strength; ASPIRIN® 81mg; ASPIRIN® 81mg Quick Chews. Date of revision: September 16, 2014.
- Boston Collaborative Drug Surveillance Group. Regular Aspirin intake and acute myocardial infarction. *Br Med J* 1974; 1: 440-3.
- Bousser MG, Eschwege E, Haguenu M, Lefauconnier JM, Thibult N, et al. "AICLA" controlled trial of Aspirin and dipyridamole in the secondary prevention of atherothrombotic cerebral ischemia. *Stroke* 1983; 14:5-14.
- Boyd EM. Analgesic abuse. Maximal tolerated daily doses of acetylsalicylic acid. *Can Med Assoc J* 1968; 99: 790-8.
- Boyd EM. The acute oral toxicity of acetylsalicylic acid. *Toxic Appl Pharmac* 1959; 1: 229-39.
- Breiddin K, Loew D, Lechner K, Oberla K, Walter E. The German_Austrian trial. A comparison of acetylsalicylic acid, placebo and phenprocoumon in secondary prevention of myocardial infarction. *Circulation* 1980; 62 (Suppl V): V63-V72.
- Breiddin K, Loew D, Lechner K, Oberla K, Walter E. Secondary prevention of myocardial infarction: Comparison of treatment with acetylsalicylic acid, phenprocoumon or placebo. A multicentre 2 year prospective study. *Int*

- Congr Ser 1979; 470: 263-8.
- Cairns JA, Gent M, Singer J, Finnie KJ, Froggatt GM, et al. Aspirin, sulfinpyrazone, or both in unstable angina. Results of a Canadian multicenter trial. *N Engl J Med* 1985; 313: 1369-75.
- Canadian Cooperative Study Group. A randomized trial of Aspirin and sulfinpyrazone in threatened stroke. *N Engl J Med* 1978; 299: 53-9.
- Christopher J. Needs and Peter M. Brooks. Clinical Pharmacokinetics of the Salicylates. *Clinical Pharmacokinetics* 1985 10: 164-177.
- Clive DM, Stoff JS. Renal syndromes associated with non-steroidal anti-inflammatory use. *N Engl J Med*. 1984 Mar 1;310(9):563-72.
- Clopath P. The effect of acetylsalicylic acid (ASA) on the development of atherosclerotic lesions in miniature swine. *Br J Exp Path* 1980; 61: 440-3.
- Coronary Drug Project Research Group. Aspirin in coronary heart disease. *J Chron Dis* 1976; 29: 625-42.
- Coronary Drug Project Research Group. Aspirin in coronary heart disease. *Circulation* 1980; 62 (Suppl V): V59-V62.
- Craven LL. Acetylsalicylic acid, Possible preventive coronary thrombosis. *Ann West Med Surg* 1950; 4: 95-9.
- Craven LL. Prevention of coronary and cerebral thrombosis. *Miss Valley Med J* 1956; 78: 213-5.
- Catella-Lawson F, Reilly MP, Kapoor SC, Cucchiara AJ, DeMarco S, Tournier B, Vyas SN, FitzGerald GA. Cyclooxygenase inhibitors and the antiplatelet effects of Aspirin. *N Engl J Med*. 2001 Dec 20;345(25):1809-17.
- Czaplicki S, Gietka J, Suzek K. The frequency of coronary heart disease and myocardial infarction in rheumatoid arthritis patients. *Cor Vasa* 1978; 20: 249-54.
- Dalton SO, Johansen C, Mellekjær L, Nørgård B, Sørensen HT, Olsen JH. Use of selective serotonin reuptake inhibitors and risk of upper gastrointestinal tract bleeding: a population- based cohort study. *Arch Intern Med*. 2003 Jan 13;163(1):59-64.
- Danese CA, Voletti CD, Weiss HJ. Protection by Aspirin against experimentally induced arterial thrombosis in dogs. *Thrombos Diathes Haemorrh* 1971; 25: 288-96.
- Dargan PI, Wallace CI, and Jones AL. An evidence based flowchart to guide the management of acute salicylate (Aspirin) overdose. *Emer Med J* 2002;19:206-209.
- De Caterina R, Giannessi D, Boem A, Bernini W, Battaglia D, Michelassi C, Dell'Amico F, L'Abbate A, Patrignani P, Patrono C. Equal antiplatelet effects of Aspirin 50 or 324 mg/day in patients after acute myocardial infarction. *Thromb Haemostas* 1985; 54: 528-32.
- De Gaetano G. Primary prevention of vascular disease by ASPIRIN. *Lancet* 1988; 1: 1093-1094.
- De Swiet M, Fryers G. Review: the use of aspirin in pregnancy. *J Obstet Gynecol*. 1990;10:467-482.
- Dunn MJ, Scharschmidt L, Zambraski E. Mechanisms of nephrotoxicity of non-steroidal anti- inflammatory drugs. *Arch Toxicol Suppl*. 1984;7:328-37.
- Dunn MJ, Zambraski EJ. Renal effects of drugs that inhibit prostaglandin synthesis. *Kidney Int*.1980 Nov;18(5):609-622
- Editorial. Aspirin after myocardial infarction. *Lancet* 1980; 1: 1172-3.
- Editorial. Trials of drugs for prevention of 'secondary' MIs indecisive. *Hosp Prac (April)* 1980: 53-4.
- Elwood PC, Sweetnam PM. Aspirin and secondary mortality after myocardial infarction. *Lancet* 1979; 2: 1313-5.
- Evans M, Ford CM, Bellocco R, Fitzmaurice G, Fryzek JP, McLaughlin JK, Nyrén O, Elinder CG. Acetaminophen, aspirin and progression of advanced chronic kidney disease. *Nephrol Dial Transplant*. 2009

Jun;24(6):1908-18.

Farah AE, Rosenberg F. Potential therapeutic application of Aspirin and other cyclo-oxygenase inhibitors. *Br J Clin Pharmacol* 1980; 10: 261S-78S.

Fields WS, Lemak NA, Frankowski RF, Hardy RJ. Controlled trial of Aspirin in cerebral ischemia. *Stroke* 1977; 8: 301-16.

Fitzgerald GA, Oates JA, Hawiger J, Maas RL, Roberts LJ 2nd, Lawson JA and Brash AR. Endogenous biosynthesis of prostacyclin and thromboxane and platelet function during chronic administration of Aspirin in man. *Clin Invest* 1983; 71: 676-88.

Flower RJ, Moncada S, Vane JR. Analgesic-antipyretics and anti-inflammatory agents; drugs employed in the treatment of gout. In: *The Pharmacological Basis of Therapeutics*, Chapter 29, 1980, 682-692.

Fored CM, Ejerblad E, Lindblad P, Fryzek JP, Dickman PW, Signorello LB, Lipworth L, Elinder CG, Blot WJ, McLaughlin JK, Zack MM, Nyrén O. Acetaminophen, aspirin, and chronic renal failure. *N Engl J Med*. 2001 Dec 20;345(25):1801-8.

Genton E. A perspective on platelet - suppressant drug treatment in coronary artery and cerebrovascular disease. *Circulation* 1980; 62: V111-V121.

Glader BE. Evaluation of the hemolytic role of aspirin in glucose-6-phosphate dehydrogenase deficiency. *J Pediatr*. 1976 Dec;89(6):1027-8.

Green LH, Seropian E, Handin RI. Platelet activation during exercise-induced myocardial ischemia. *N Engl J Med* 1980; 302: 193-7.

Grosser N, Schröder H. Aspirin protects endothelial cells from oxidant damage via the nitric oxide-cGMP pathway. *Arterioscler Thromb Vasc Biol*. 2003 Aug 1;23(8):1345-51. Epub 2003 Jun 26.

Guslandi M. Gastric toxicity of antiplatelet therapy with low-dose Aspirin. *Drugs* 1997; 53: 1-5.

Haddad LM et al. *Clinical Management of Poisoning and Drug Overdose* 3rd ed. Chap 50, Salicylate Toxicity. W.B. Saunders Company ©1998, p. 675-687.

Haft JJ. Platelets and coronary artery disease. *Prim Card* 1979; June: 97-104.

Hansson L, Zanchetti A, Carruthers SG, Dahlöf B, Elmfeldt D, Julius S, Ménard J, Rahn KH, Wedel H, and Westerling S. Effects of intensive blood-pressure lowering and low-dose Aspirin in patients with hypertension: principal results of the Hypertension Optimal Treatment (HOT) randomised trial. HOT Study Group. *Lancet*. 1998 Jun 13;351(9118):1755-62.

Harker LA, Slichter SJ. Studies of platelet and fibrinogen kinetics in patients with prosthetic heart valves. *N Engl J Med* 1970; 283: 1302-5.

Harris WH, Salzman EW, Athanasoulis CA, Waltman AC, Baum S, De Sanctis RW. Aspirin prophylaxis of venous thromboembolism after total hip replacement. *N Engl J Med* 1977; 297: 1246-9.

Harris WH, Salzman EW, Athanasoulis CA, Waltman AC, Baum S, De Sanctis RW. Comparison of warfarin, low molecular weight dextran, Aspirin and subcutaneous heparin in prevention of venous thromboembolism following total hip replacement. *J Bone Joint Surg* 1974; 56: 1552-62.

Harrison MJG, Marshall J, Meadows JC, Russell RWR. Effect of Aspirin in Amaurosis Fugax. *Lancet* 1971; 2: 743-4.

Grosser N, Abate A, Oberle S, Vreman HJ, Dennery PA, Becker JC, Pohle T, Seidman DS, Schröder H. Heme oxygenase-1 induction may explain the antioxidant profile of Aspirin. *Biochem Biophys Res Commun*. 2003 Sep 5;308(4):956-60.

Hennekens CH, Peto R, Hutchison GB, Doll R. An overview of the British and American Aspirin studies. *N Engl J Med* 1988; 318: 923-4.

- Hirsh J, Dalen JE, Fuster V, Harker LB, Salzman EW. Aspirin and other platelet-active drugs. The relationship between dose, effectiveness and side effects. *Chest* 1992; 102: 327S-36S.
- Hoffman W, Forster W. Two year Cottbus reinfarction study with 30 mg ASA per day. Prostaglandins, Leukotrienes and Essential Fatty Acid 1991; 44: 159-69.
- Hoffman W, Nitschke M, Muche J, Kampe W, Handreg W, Forster W. Reevaluation of the Cottbus reinfarction study with 30 mg Aspirin per day 4 years after the end of the study. Prostaglandins, Leukotrienes and Essential Fatty Acids 1991; 42: 137-9.
- Hogben AMC, Tocco DJ, Brodie BB, Schanker LS. On the mechanism of intestinal absorption of drugs. *J Pharm Ther* 1959; 125: 275-82.
- Hollifield, JW. Failure of Aspirin to Antagonize the Antihypertensive Effect of Spironolactone in Low-Renin Hypertension. *Southern Medical Journal* 1976; 69(8): 1034-1036.
- Hsia J, Hamilton WP, Kleiman N, Roberts R, Chaitman BR, Ross AM for the Heparin- Aspirin Reperfusion Trial (HART) Investigators. A comparison between heparin and low-dose Aspirin as adjunctive therapy with tissue plasminogen activator for acute myocardial infarction. *N Engl J Med* 1990; 323: 1433-7.
- Hume M, Bierbaum B, Kurlakose TX, Surprenant J. Prevention of post-operative thrombosis by ASPIRIN. *Amer J Surg* 1977; 133: 420-2.
- ISIS-2 (Second International Study of Infarct Survival) Collaborative Group. Randomised trial of intravenous streptokinase, oral Aspirin, both, or neither among 17,187 cases of suspected acute myocardial infarction: ISIS-2. *Lancet*. 1988 Aug 13;2(8607):349-60.
- Jakubowski JA, Stampfer MJ, Vaillancourt R, Deykin D. Cumulative antiplatelet effect of low- dose enteric coated ASPIRIN. *Br J Haematol* 1985; 60: 635-42.
- James MJ, Walsh JA. Effects of aspirin and alcohol on platelet thromboxane synthesis and vascular prostacyclin synthesis. *Thromb Res*. 1985 Sep 1;39(5):587-93.
- Jennings JJ, Harris WH, Sarmiento A. A clinical evaluation of Aspirin prophylaxis of thromboembolic disease after total hip arthroplasty. *J Bone Joint Surg* 1976; 58: 926-8.
- Juul-Moller S, Edvardsson N, Jahnmatz B, Rosen A, Sorensen S, Omblus R, for the Swedish Angina Pectoris Aspirin Trial (SAPAT) Group: Double-blind trial of Aspirin in primary prevention of myocardial infarction in patients with stable chronic angina pectoris. *Lancet* 1992; 340: 1421-5.
- Kaufman DW, Kelly JP, Wiholm BE, Laszlo A, Sheehan JE, Koff RS, Shapiro S. The risk of acute major upper gastrointestinal bleeding among users of aspirin and ibuprofen at various levels of alcohol consumption. *Am J Gastroenterol*. 1999 Nov;94(11):3189-3196.
- Kumpuris AG, Luchi RJ, Waddell CC, Miller RR. Production of circulating platelet aggregates by exercise in coronary patients. *Circulation* 1980; 61: 62-5.
- Kurth T, Glynn RJ, Walker AM, Rexrode KM, Buring JE, Stampfer MJ, Hennekens CH, Gaziano JM. Analgesic use and change in kidney function in apparently healthy men. *Am J Kidney Dis*. 2003 Aug;42(2):234-44.
- Levin A, Stevens L, McCullough PA. Cardiovascular disease and the kidney: tracking a killer in chronic kidney disease. *Postgrad Med*. 2002 Apr;111(4):53-60.
- Levin A. Clinical epidemiology of cardiovascular disease in chronic kidney disease prior to dialysis. *Semin Dial*. 2003 Mar-Apr;16(2):101-5.
- Lewis HD Jr, Davis JW, Archibald DG, Steinke WE, Smitherman TC, et al. Protective effects of Aspirin against acute myocardial infarction and death in men with unstable angina. Results of a Veterans Administration Cooperative Study. *N Engl J Med* 1983; 309: 396-403.
- Lipton RB, Goldstein J, Baggish JS, Yataco AR, Sorrentino JV, Quiring JN. Aspirin is efficacious for the treatment of acute migraine. *Headache*. 2005 Apr;45(4):283-92.

- Lindblad B, et al. Does low-dose acetylsalicylic acid prevent stroke after carotid surgery? A double-blind, placebo-controlled randomized trial. *Stroke* 1993; 24:1125-8.
- MacDonald TM, Wei L. Effect of ibuprofen on cardioprotective effect of Aspirin. *Lancet*. 2003 Feb 15;361(9357):573-4.
- Malseed R, Malseed Z. Aspirin: a pharmacologic profile. *Amer J Pharm* 1978; July-Aug: 150: 99-106.
- Mangano DT.. Aspirin and mortality from coronary bypass surgery. *N Engl J Med*. 2002 Oct 24;347(17):1309-17.
- McCann RL, Hagen P, Fuchs JCA. Aspirin and dipyridamole decrease intimal hyperplasia in experimental vein grafts. *Ann Surg* 1980; 191: 238-43.
- McKenna R, Bachmann F, Kaushal SP, Galante JO. Thromboembolic disease in patients undergoing total knee replacement. *J Bone Joint Surg* 1976; 58: 928-32.
- Medical Research Council's General Practice Research Framework. Thrombosis prevention trial:randomized trial of low-intensity oral anticoagulation with warfarin and low-dose Aspirin in the primary prevention of ischaemic heart disease in men at increased risk. *Lancet* 1998;351:233-41.
- Mehta J, Mehta P, Burger C, Pepine CJ. Platelet function studies in coronary heart disease. IV. Effect of Aspirin. *Atherosclerosis* 1978; 31: 169-75.
- Mehta J, Mehta P, Pepine CJ, Contic CR. Platelet aggregation studies in coronary heart disease. VII. Effect of Aspirin and tachycardia stress on aortic and coronary venous blood. *Am J Cardiol* 1980; 45: 945-51.
- Mehta P, Mehta J. Platelet function studies in coronary heart disease. V. Evidence for enhanced platelet microthrombus formation activity in acute myocardial infarction. *Am J Cardiol* 1979; 43: 757-60.
- Montgomery PR, Berger LG, Mitenko PA, Sitar DS. Salicylate metabolism: effects of age and sex in adults. *Clin Pharmacol Ther*. 1986 May;39(5):571-6.
- Morley J. Mechanism of action of Aspirin in inflammation. *Proc Roy Soc Med* 1977; 70: 326. Moschos CB, Haider B, De La Cruz C, Lyons MM, Regan TJ. Antiarrhythmic effects of ASPIRIN during non_thrombotic coronary occlusion. *Circulation* 1978; 57:681-4.
- Mundall J, Quintero P, Von Kaulla KN, Harmon R, Austin J. Transient monocular blindness and increased platelet aggregability treated with ASA. *Neurology* 1972; 22: 280-5.
- Murray MD, Black PK, Kuzmik DD, Haag KM, Manatunga AK, Mullin MA, Hall SD, Brater DC. Acute and chronic effects of nonsteroidal antiinflammatory drugs on glomerular filtration rate in elderly patients. *Am J Med Sci*. 1995 Nov;310(5):188-97.
- Murray MD, Brater DC. Effects of NSAIDS on the kidney. *Prog Drug Res*. 1997;49:155-71. Orme M. Aspirin all round? *Br Med J* 1988; 296: 307-8.
- Orr J, Abbott F, Farrell K, Ferguson S, Sheppard I, and W Godolphin, Interaction between valproic acid and Aspirin in epileptic children: Serum protein binding and metabolic effects. *Clin. Pharmacol.Ther*. 1982;31:642-649.
- Packham MA, Mustard JF. Pharmacology of platelet affecting drugs. *Circulation* 1980; 62: V26-V41.
- Patrignani P, Filabozzi P, Patrono C. Selective cumulative inhibition of platelet thromboxane production by low-dose Aspirin in healthy subjects. *J Clin Invest* 1982; 69: 1366-72.
- Persantine_ASPIRIN Reinfarction Study (PARIS) Research Group: The Persantine Aspirin reinfarction study. *Circulation* 1980; 62 (Suppl V): V85-V88.
- Pick R, Chediak J, Glick G. ASPIRIN inhibits development of coronary atherosclerosis in cynomolgus monkeys (*Macaca Fascicularis*) fed on atherogenic diet. *J Clin Invest* 1970; 63: 158-62.
- Pugliese F, Ciabattoni G. The role of prostaglandins in the control of renal function: renal effects of nonsteroidal anti-inflammatory drugs. *Clin Exp Rheumatol*. 1984 Oct-Dec;2(4):345- 52.

- Relman AS. Aspirin for the primary prevention of myocardial infarction. *N Engl J Med* 1988; 318: 245-6.
- Renaud S, Godu J. Thrombosis prevention by acetylsalicylic acid in hyperlipemic rats. *CMAJ* 1970; 103: 1037-40.
- RISC Group. Risk of myocardial infarction and death during treatment with low dose Aspirin and intravenous heparin in men with unstable coronary artery disease. *Lancet* 1990; 336: 827-30.
- Roberts MS, Joyce RM, McLeod LJ, Vial JH, Seville PR. Slow-release Aspirin and prostaglandin inhibition. *Lancet* 1986; 1(8490): 1153-4.
- Ross R, Glomset JA. Pathogenesis of atherosclerosis. *N Engl J Med* 1976; 295: 369-377, 420-5.
- Roth GJ, Stanford N, Majenus PW. Acetylation of prostaglandin synthase by Aspirin. *Proc Nat Acad Sci* 1975; 72: 3073-6.
- Rowland M, Riegelman S. Pharmacokinetics of acetylsalicylic acid and salicylic acid after intravenous administration in man. *J Pharm Sci* 1968; 57: 1313-9.
- Salicylate metabolism: effects of age and sex in adults. *Clin Pharmacol Ther.* 1986 May;39(5):571-6.
- SALT Collaborative Group. Swedish Aspirin Low-Dose Trial (SALT) of 75 mg Aspirin as secondary prophylaxis after cerebrovascular ischaemic events. *Lancet* 1991; 338: 1345-9.
- Salzman EW, Harris WH, De Sanctis RW. Reduction in venous thromboembolism by agents affecting platelet function. *N Engl J Med* 1971; 284: 1287-92.
- Sarnak MJ, Levey AS. Cardiovascular disease and chronic renal disease: a new paradigm. *Am J Kidney Dis.* 2000 Apr;35(4 Suppl 1): S117-31.
- Schafer A, Handin R. The role of platelets in thrombotic and vascular disease. *Proj Cardio Dis* 1979; 22: 31-52.
- Schrör K. Acetylsalicylic acid. Chapter 3 Toxicity and Drug Safety: Section 3.2.2.2 Mode of Aspirin Action; WILEY-VCH Verlag GmbH & Co. KGaA Weinheim 2009; 180-181.
- Shahidi NT, Westring DW. Acetylsalicylic acid-induced hemolysis and its mechanism. *J Clin Invest.* 1970 Jul;49(7):1334-40.
- Shalev O. Long-term, low-dose aspirin is safe in glucose-6-phosphate dehydrogenase deficiency. *DICP.* 1991 Oct;25(10):1074-75.
- Slone D, Siskind V, Heinonen OP, Monson RR, Kaufman DW, Shapiro S. Aspirin and congenital malformations. *Lancet.* 1976 Jun 26;1(7974):1373-5.
- Smith MJH. Plasma-salicylate concentrations after small doses of acetylsalicylic acid. *J Pharm Pharmacol* 1951; 3: 409-14.
- Soreff J, Johnson H, Diener L, Göransson L. Acetylsalicylic acid in a trial to diminish thromboembolic complications after elective hip surgery. *Acta Orthop Scand* 1975; 46: 246-55.
- Steering Committee of the Physicians' Health Study Research Group. Final report on the Aspirin component of the ongoing Physicians' Health Study. Steering Committee of the Physicians' Health Study Research Group. *N Engl J Med.* 1989 Jul 20;321(3):129-35.
- Stockman III JA, Lubin B, Oski FA. Aspirin-induced hemolysis: the role of concomitant oxidant (H₂O₂) challenge. *Pediatr Res.* 1978 Sept;12(9):927-31.
- Tata LJ, Fortun PJ, Hubbard RB, Smeeth L, Hawkey CJ, Smiths CJP, Whitaker HJ, Farrington CP, Card TR, West J. Does concurrent prescription of selective serotonin reuptake inhibitors and non-steroidal anti-inflammatory drugs substantially increase the risk of upper gastrointestinal bleeding? *Aliment Pharmacol Ther.* 2005 Aug 1;22(3):175-81.
- The Dutch TIA Trial Study Group. A comparison of two doses of ASPIRIN (30 mg vs. 283 mg a day) in patients

after a transient ischemic attack or minor ischemic stroke. N Engl J Med 1991; 325: 1261-6.

UK_TIA Study Group. United Kingdom transient ischaemic attack (UK-TIA) Aspirin trial: interim results. Br Med J 1988; 296: 316-20.

Vane JR. Inhibition of prostaglandin synthesis as a mechanism of action for Aspirin-like drugs. Nature New Biol 1971; 231: 232-5.

Verheugt FWA, van der Loarse A, Funke-Kupper AJ, Sterkman LGW, Galema TW, Roos JP. Effects of early intervention with low-dose Aspirin (100 mg) on infarct size, reinfarction and mortality in anterior wall acute myocardial infarction. Am J Cardiol 1990; 66: 267-70.

Wallentin LC and The Research Group on Instability in Coronary Artery Disease in Southeast Sweden. Aspirin (75 mg/day) after an episode of unstable coronary artery disease: Long-term effects on the risk for myocardial infarction, occurrence of severe angina and the need for revascularization. J Am Coll Cardiol 1991; 18: 1587-93.

Weiss HJ, Aledort LM, Kochwa S. The effect of salicylates on the hemostatic properties of platelets in man. J Clin Invest 1968; 47: 2169-80.

Weiss HJ. Antiplatelet therapy (second of two parts). N Engl J Med 1978; 298: 1403-6.

Wessinger S, Kaplan M, Choi L, Williams M, Lau C, Sharp L, Crowell MD, Keshavarzian A, Jones MP. Increased use of selective serotonic reuptake inhibitors in patients admitted with gastrointestinal hemorrhage: a multicenter retrospective analysis. Alimet Pharmacol Ther. 2006 Apr 1;23(7):934-44.

Wright HN. Chronic toxicity studies of analgesic and anti-pyretic drugs and congeners. Toxicol Appl Pharmacol 1967; 11: 280-92.

Young VP, Giles AR, Pater J, Corbett WE. Sex differences in bleeding time and blood loss in normal subjects following Aspirin ingestion. Thromb Research 1980; 20: 705-9.

Zambraski EJ, Dunn MJ. Renal effects of aspirin 1992 Edited by Vane JR and Botting RM in: Aspirin and other salicylates. London-New York-Tokyo-Melbourne-Madras: Chapman & Hall Medical; 510-530.

Zuik M, Mandel MA. Methotrexate-salicylate interaction: a clinical and experimental study. Surg Forum. 1976;26:567-9.

**PART III: CONSUMER INFORMATION ONLY
PROVIDED BY HEALTH CARE PROFESSIONALS**

RIVASA 81 mg EC

(Acetylsalicylic Acid (ASA) Delayed Release Tablets, USP)

This leaflet is part III of a three-part "Product Monograph" published when **RIVASA 81 mg EC** was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about **RIVASA 81 mg EC**. Contact your doctor or pharmacist if you have any questions about the drug. Also see package insert for additional information.

ABOUT THIS MEDICATION

What the medication is used for:

RIVASA 81 mg EC can help save your life in the following situations:

- to help prevent a first heart attack in those who are at increased risk, or
- to help prevent a second heart attack or stroke in those who have already had such an event.

RIVASA 81 mg EC may help save your life if you think you are having a heart attack.

FOR PREVENTION OF A FIRST NON-FATAL HEART ATTACK (DAILY THERAPY):

Your doctor may recommend you take **RIVASA 81 mg EC** to help reduce the risk of a first non-fatal heart attack because you are at risk of having a heart attack. There is no evidence that this product reduces the risk of a first fatal heart attack, nor first strokes (fatal and non-fatal), nor death due to any cardiovascular problems. Your doctor will assess the appropriate balance of possible benefit of this product against the potential risk of stomach bleeding and stroke. Factors that increase your risk include high blood pressure, high cholesterol, diabetes, family history of heart disease, increased age, overweight and smoking. You should follow your doctor's instructions carefully. Please notify your doctor if you intend to stop taking this medication.

USE DURING A HEART ATTACK

If you think you are having a heart attack, call 911 immediately, then chew or crush and swallow two **RIVASA 81 mg EC**. It is important to chew or crush the product, to ensure this medicine quickly works. Then get to a hospital immediately for medical attention. Inform the emergency services / hospital that you have taken **RIVASA 81 mg EC**. Taking **RIVASA 81 mg EC** at the first signs and symptoms can reduce your risk of dying from the heart attack. The signs and symptoms of a heart attack include:

- uncomfortable pressure, fullness, squeezing or pain in the centre of the chest that lasts more than a few minutes, or goes away quickly and comes back,
- pain that spreads to the shoulders, neck or arms,
- chest discomfort with lightheadedness, fainting, sweating, nausea or shortness of breath.

Chest pain is the most common symptom in both sexes, but women may also experience other symptoms such as unusual fatigue that gets worse with activity, difficulty of breathing, heartburn/nausea and /or vomiting unrelieved by antacids, tightening and pain in the chest that may extend into the neck,

jaws and shoulders, general feeling of weakness, paleness, sweating).

At the hospital, the doctor will then recommend appropriate therapy.

FOR PREVENTION OF A SECOND HEART ATTACK OR STROKE (DAILY THERAPY)

Your doctor may recommend you take **RIVASA 81 mg EC** daily to help prevent a second heart attack or stroke. After having experienced a first heart attack or stroke, you can be at increased risk of experiencing a second one. You may also be at risk for heart disease and stroke because you may be overweight, a smoker, have an inactive lifestyle, high blood pressure, are under stress or have high blood cholesterol.

Following your doctor's instructions concerning the use of **RIVASA 81 mg EC** and the changes in diet, exercise and lifestyle he/she may have prescribed, will provide you with your best opportunity to avoid experiencing a second heart attack or stroke. Always contact your doctor if you experience any difficulties.

What it does:

RIVASA 81 mg EC belongs to a group of medicines called antiplatelet drugs. Platelets are very small structures in blood, smaller than red or white blood cells, which clump together during blood clotting. By preventing this clumping, antiplatelet drugs reduce the chances of blood clots forming (a process called thrombosis).

When it should not be used:

DO NOT TAKE if you:

- are allergic to ASA, salicylates, non-steroidal anti-inflammatory drugs (NSAIDs)/pain relievers/fever reducers, or other ingredients in the product
- have an ulcer, history of ulcers or are prone to bleeding
- have active or severe liver or kidney disease or congestive heart failure
- have a history of asthma caused by salicylates or other NSAIDs
- are using methotrexate at doses of 15 mg/week or more
- are in the last trimester of pregnancy because it may cause problems in the unborn child or complications during delivery

What the medicinal ingredient is:

acetylsalicylic acid (ASA)

What the important nonmedicinal ingredients are:

Colloidal silica, corn starch, FD&C blue #1, hypromellose, methacrylic acid copolymer, microcrystalline cellulose, purified water, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate.

What dosage forms it comes in:

RIVASA 81 mg EC comes as enteric-coated tablets.

WARNINGS AND PRECAUTIONS

Your doctor will have asked you many questions about your health, lifestyle, and medications before recommending RIVASA 81 mg EC. That is why it is very important that you tell your doctor all such information. If you have forgotten to tell your doctor about any of the following, call your doctor or pharmacist before you take this medicine (or any medicine):

- asthma, high blood pressure, heart disease, gout or other serious conditions
- age 60 years or older
- stomach problems such as heartburn
- **STOMACH BLEEDING WARNING:** contains a NSAID which **may cause severe stomach bleeding**
- impaired liver/kidney or impaired cardiovascular circulation (renal vascular disease, congestive heart failure, volume depletion, major surgery, sepsis or major hemorrhagic events);
- history of blood clotting defects;
- have severe anemia;
- have severe glucose-6-phosphate dehydrogenase (G6PD) deficiency;
- are trying to conceive, pregnant or breast-feeding or
- will be having surgery in five to seven days

● **CAUTION:** Contains enough drug to seriously harm a child; **KEEP OUT OF THEIR REACH. DO NOT GIVE** to children/teenagers less than 18 years of age who have chicken pox or cold/flu symptoms before a doctor is consulted about Reye's Syndrome, a rare but serious illness reported to be associated with ASA.

INTERACTIONS WITH THIS MEDICATION

Tell your doctor if you are taking any prescription or non-prescription drugs including blood thinners, acetaminophen, anticonvulsants, anti-diabetic/arthritis/gout medicines, digoxin, glucocorticoids, methotrexate, selective-serotonin re-uptake inhibitors (a type of antidepressant), diuretics, ACE inhibitors (medication for high blood pressure), or are having 3 or more alcoholic drinks per day.

Ibuprofen may interfere with the heart protective benefits of ASPIRIN. Patients should talk to their doctor if they are on an ASPIRIN regimen and take ibuprofen for pain.

PROPER USE OF THIS MEDICATION

Usual dose:

DIRECTIONS (Adults ≥ 18 years): During a heart attack: Call 911, then chew or crush 2 – RIVASA 81 mg EC tablets.

For prevention of a first heart attack or for the prevention of a second heart attack or stroke: Take 1 to 4 tablets daily, depending on your doctor's instructions. You should take this medicine at the same time every day. This will help you to remember to take your medication. For maximum effectiveness, it is very important to take RIVASA 81 mg EC *every day* as directed by your doctor. Do not take more tablets than your doctor recommends. Your doctor may tell you to take RIVASA 81 mg EC with other medications; he or she may also tell you to eat special foods, exercise or take other steps to safeguard your health.

For daily therapy RIVASA 81 mg EC, tablets should be swallowed whole for the medicine to work properly. RIVASA 81 mg EC tablets have a special coating, *enteric coating*, which allows the tablets to pass undissolved through the stomach and on into the intestine. By dissolving in the intestine rather than the stomach, the risk of stomach upset is reduced in those with a sensitive stomach. Therefore, to maintain this protection, the tablets should not be crushed or broken.

Can I Continue to Take RIVASA 81 mg EC for Relief of Headache, Fever or Arthritis Pain?

RIVASA 81 mg EC is specially designed to reduce your risk of dying during a heart attack, to help prevent a first heart attack in those who are at increased risk and to help prevent a second heart attack or stroke. It is a smaller dose than you would need to take for a headache or other types of pain and is unlike other pain reliever products such as acetaminophen or NSAIDs e.g., ibuprofen, naproxen. Ask your doctor or pharmacist about other ASA products available (or other pain relievers such as acetaminophen or ibuprofen, naproxen or salicylates) and the correct dosage for the relief of your headache, fever or arthritic pain. Always consult with your doctor or pharmacist before taking other medications.

Overdose:

If you think you have taken too much RIVASA 81 mg EC, contact your healthcare professional, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

Missed Dose:

If you forget to take your medication, take it when you remember. But do not take *extra* medication to compensate for a missed dosage unless instructed by your doctor.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Like all medicines, RIVASA 81 mg EC may occasionally produce unwanted side effects. You should call your doctor if you experience any of the following: nausea, vomiting; stomach irritation, or pain; if you notice that you are 'bruising' more easily than you were before starting a daily dose of RIVASA 81 mg EC. Regular daily use of alcohol while on RIVASA 81 mg EC daily therapy may increase your risk of developing gastrointestinal bleeding. This is not a complete list of side effects. For any unexpected effects while taking RIVASA 81 mg EC, contact your doctor or pharmacist.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Stop use and call your doctor if you experience an allergic reaction (skin rash, hives, itching, swelling of eyes, face, lips, tongue, or throat, wheezing or breathing difficulties); stomach bleeding (feel faint, bloody vomit, vomit that looks like coffee grounds, bright red blood in stools, black or tarry stools, stomach pain that does not get better); loss of hearing, ringing or buzzing in ears or bleeding.

HOW TO STORE IT

Keep out of the site and reach of children.

Store at room temperature, 15-30°C.

Reporting Side Effects

You can help improve the safe use of health products for Canadians by reporting serious and unexpected side effects to Health Canada. Your report may help to identify new side effects and change the product safety information.

3 ways to report:

- Online at MedEffect;
- By calling 1-866-234-2345 (toll-free);
- By completing a Consumer Side Effect Reporting Form and sending it by:
 - Fax to 1-866-678-6789 (toll-free), or
 - Mail to: Canada Vigilance Program
Health Canada, Postal Locator 0701E
Ottawa, ON
K1A 0K9

Postage paid labels and the Consumer Side Effect Reporting Form are available at MedEffect.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION**If you want more information about RIVASA 81 mg EC:**

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website; the manufacturer's website www.labriva.com or by calling 1-800-363-7988.

This leaflet was prepared by:

Laboratoire Riva Inc.

660 Industriel Blvd
Blainville, Quebec, Canada
J7C 3V4

Last revised: December 21, 2015

PART III: CONSUMER INFORMATION ONLY

RIVASA 81 mg EC

(Acetylsalicylic Acid (ASA) Delayed Release Tablets, USP)

This leaflet is part III of a three-part "Product Monograph" published when **RIVASA 81 mg EC** was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about **RIVASA 81 mg EC**. Contact your doctor or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATION

What the medication is used for:

- RIVASA 81 mg EC is for doctor supervised long-term preventative therapy.
- RIVASA 81 mg EC may help save your life if you think you are having a heart attack.

What it does:

RIVASA 81 mg EC is for doctor supervised long-term preventive therapy.

USE DURING A HEART ATTACK

If you think you are having a heart attack, call 911 immediately then, chew or crush and swallow two RIVASA 81 mg EC tablets. It is important to chew or crush the product, to ensure this medicine works quickly. Then get to a hospital immediately for medical attention. Inform the emergency services / hospital that you have taken RIVASA 81 mg EC. Taking RIVASA 81 mg EC, at the first signs and symptoms can reduce your risk of dying from a heart attack.

The signs and symptoms of a heart attack include:

- uncomfortable pressure, fullness, squeezing or pain in the centre of the chest that lasts more than a few minutes, or goes away quickly and comes back,
- pain that spreads to the shoulders, neck or arms,
- chest discomfort with light-headedness, fainting, sweating, nausea or shortness of breath.

Chest pain is the most common symptom in both sexes, but women may also experience other symptoms such as unusual fatigue that gets worse with activity, difficulty of breathing, heartburn/nausea and /or vomiting unrelieved by antacids, tightening and pain in the chest that may extend into the neck, jaws and shoulders, general feeling of weakness, paleness, sweating)

At the hospital, the doctor will then recommend appropriate therapy.

When it should not be used:

DO NOT TAKE if you:

- are allergic to ASA, salicylates, non-steroidal anti-inflammatory drugs (NSAIDs)/pain relievers/fever reducers or other ingredients in the product
- have an ulcer, history of ulcers or are prone to bleeding

- have active or severe liver or kidney disease or congestive heart failure
- have a history of asthma caused by salicylates or other NSAIDs
- are using methotrexate at doses of 15 mg/week or more
- are in the last trimester of pregnancy because it may cause problems in the unborn child or complications during delivery

What the medicinal ingredient is:

acetylsalicylic acid (ASA)

What the important nonmedicinal ingredients are:

Colloidal silica, corn starch, FD&C blue #1, hypromellose, methacrylic acid copolymer, microcrystalline cellulose, purified water, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate.

What dosage forms it comes in:

RIVASA 81 mg EC comes as enteric-coated tablets.

WARNINGS AND PRECAUTIONS

Your doctor will have asked you many questions about your health, lifestyle, and medications before recommending RIVASA 81 mg EC. That is why it is very important that you tell your doctor all such information. If you have forgotten to tell your doctor about any of the following, call your doctor or pharmacist before you take this medicine (or any medicine):

- asthma, high blood pressure, heart disease, gout or other serious conditions
- age 60 years or older
- stomach problems such as heartburn
- **STOMACH BLEEDING WARNING:** contains a NSAID which may cause severe stomach bleeding
- impaired liver/kidney or impaired cardiovascular circulation (renal vascular disease, congestive heart failure, volume depletion, major surgery, sepsis or major hemorrhagic events);
- history of blood clotting defects;
- severe anemia;
- severe glucose-6-phosphate dehydrogenase (G6PD) deficiency;
- are trying to conceive, pregnant or breast-feeding or
- will be having surgery in five to seven days

● **CAUTION:** Contains enough drug to seriously harm a child; **KEEP OUT OF THEIR REACH. DO NOT GIVE** to children/teenagers less than 18 years of age who have chicken pox or cold/flu symptoms before a doctor is consulted about Reye's Syndrome, a rare serious illness reported to be associated with ASA.

INTERACTIONS WITH THIS MEDICATION

Tell your doctor if you are taking any prescription or nonprescription drugs including blood thinners, acetaminophen, anticonvulsants, anti-diabetic/arthritis/gout medicines, digoxin, glucocorticoids, methotrexate, selective serotonin re-uptake inhibitors (a type of antidepressants), diuretics, ACE inhibitors (medication for high blood pressure), or if you are having 3 or more alcoholic drinks per day. Do not use ibuprofen if you are taking RIVASA 81 mg EC for preventative therapy without talking to a doctor or pharmacist as ibuprofen may interfere with the preventive benefits of RIVASA 81 mg EC.

PROPER USE OF THIS MEDICATION

Usual dose:

DIRECTIONS (Adults ≥ 18 years): For doctor supervised long-term preventative therapy: Take 1 to 4 tablets daily, depending on your doctor's instructions. You should take this medicine at the same time every day. This will help you to remember to take your medication. For maximum effectiveness, it is very important to take RIVASA 81 mg EC *every day* as directed by your doctor. Do not take more tablets than your doctor recommends. Your doctor may tell you to take RIVASA 81 mg EC with other medications; he or she may also tell you to eat special foods, exercise or take other steps to safeguard your health. RIVASA 81 mg EC tablets have a special coating, *enteric coating*, which allows them to pass undissolved through the stomach and into the intestine. By dissolving in the intestine rather than the stomach, the risk of stomach upset is reduced. Therefore, to maintain this protection, the tablets should not be crushed or broken.

During a heart attack: Call 911, then, crush or chew 2 tablets.

Can I continue to take RIVASA 81 mg EC for relief of headache, fever or arthritis pain?

RIVASA 81 mg EC is specially designed for doctor supervised long-term preventative therapy. It is a smaller dose than you would need to take for a headache or other types of pain and is unlike other pain reliever products such as acetaminophen or NSAIDS e.g., ibuprofen, naproxen. Ask your doctor or pharmacist about other acetylsalicylic acid products available (or other pain relievers such as acetaminophen, ibuprofen, naproxen or salicylates) and the correct dosage for the relief of your headache, fever or arthritic pain. Always consult with your doctor or pharmacist before taking other medications.

Overdose:

If you think you have taken too much RIVASA 81 mg EC, contact your healthcare professional, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

Missed Dose:

If you forget to take your medication, take it when you remember.

But do not take *extra* medication to compensate for a missed dosage unless instructed by your doctor.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Like all medicines, RIVASA 81 mg EC may occasionally produce unwanted side effects. You should call your doctor if you experience: nausea, vomiting; stomach irritation, or pain; if you notice that you are 'bruising' more easily than you were before starting a daily dose of RIVASA 81 mg EC. Regular daily use of alcohol while on RIVASA 81 mg EC daily therapy may increase your risk of developing gastrointestinal bleeding. This is not a complete list of side effects. For any unexpected effects while taking RIVASA 81 mg EC, contact your doctor or pharmacist.

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Stop use and call your doctor if you experience an allergic reaction (skin rash, hives, itching, swelling of eyes, face, lips, tongue, or throat, wheezing or breathing difficulties); stomach bleeding (feel faint, bloody vomit, vomit that looks like coffee grounds, bright red blood in stools, black or tarry stools, stomach pain that does not get better); loss of hearing, ringing or buzzing in ears or bleeding.

HOW TO STORE IT

Keep out of the sight and reach of children.
Store at room temperature, 15-30°C.

Reporting Side Effects

You can help improve the safe use of health products for Canadians by reporting serious and unexpected side effects to Health Canada. Your report may help to identify new side effects and change the product safety information.

3 ways to report:

- Online at MedEffect;
- By calling 1-866-234-2345 (toll-free);
- By completing a Consumer Side Effect Reporting Form and sending it by:
 - Fax to 1-866-678-6789 (toll-free), or
 - Mail to: Canada Vigilance Program
Health Canada, Postal Locator 0701E
Ottawa, ON
K1A 0K9

Postage paid labels and the Consumer Side Effect Reporting Form are available at MedEffect.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

If you want more information about RIVASA 81 mg EC:

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website; the manufacturer's website www.labriva.com or by calling 1-800-363-7988.

This leaflet was prepared by:

Laboratoire Riva Inc.
660 Industriel Blvd
Blainville, Quebec, Canada
J7C 3V4

Last revised: December 21, 2015