

PRODUCT MONOGRAPH

PrRIVA-RANITIDINE
Ranitidine Tablets, USP
150 mg and 300 mg

Histamine H₂-receptor antagonist

LABORATOIRE RIVA INC.
660 Boul. Industriel
Blainville, Quebec
J7C 3V4

Date of Revision:
February 3, 2017

www.labriva.com

Control No.: 201549

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION	3
SUMMARY PRODUCT INFORMATION	3
INDICATIONS AND CLINICAL USE.....	3
CONTRAINdicATIONS	4
WARNINGS AND PRECAUTIONS.....	4
ADVERSE REACTIONS.....	6
DRUG INTERACTIONS	7
DOSAGE AND ADMINISTRATION.....	8
OVERDOSAGE	10
ACTION AND CLINICAL PHARMACOLOGY	10
STORAGE AND STABILITY.....	12
DOSAGE FORMS, COMPOSITION AND PACKAGING	12
PART II: SCIENTIFIC INFORMATION.....	13
PHARMACEUTICAL INFORMATION.....	13
CLINICAL TRIALS.....	14
DETAILED PHARMACOLOGY	15
TOXICOLOGY	18
REFERENCES	22
PART III: CONSUMER INFORMATION	25

PrRIVA-RANITIDINE
Ranitidine Tablets, USP
150 mg and 300 mg

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form/Strength	All Nonmedicinal Ingredients
Oral	Tablets: 150 mg and 300 mg	Colloidal Silicon Dioxide, Croscarmellose Sodium, Lecithin, Magnesium Stearate, Microcrystalline Cellulose, Polyvinyl Alcohol Part Hydrolyzed, Talc, Titanium Dioxide and Xanthan Gum.

INDICATIONS AND CLINICAL USE

RIVA-RANITIDINE (ranitidine hydrochloride) tablets are indicated for the treatment of duodenal ulcer, benign gastric ulcer, reflux esophagitis, post-operative peptic ulcer, Zollinger-Ellison Syndrome, and other conditions where reduction of gastric secretion and acid output is desirable. These include the following:

- the treatment of nonsteroidal anti-inflammatory drug (NSAID)- induced lesions, both ulcers and erosions, and their gastrointestinal (GI) symptoms and the prevention of their recurrence;
- the prophylaxis of GI haemorrhage from stress ulceration in seriously ill patients;
- the prophylaxis of recurrent haemorrhage from bleeding ulcers;
- the prevention of Acid Aspiration Syndrome from general anaesthesia in patients considered to be at risk for this, including obstetrical patients in labour, and obese patients.

In addition, RIVA-RANITIDINE is indicated for the prophylaxis and maintenance treatment of duodenal or benign gastric ulcer in patients with a history of recurrent ulceration.

Geriatrics (> 65 years of age): see WARNINGS AND PRECAUTIONS, Special Populations, Geriatrics.

Pediatrics (<18 years of age): see WARNINGS AND PRECAUTIONS, Special Populations, Pediatrics.

CONTRAINDICATIONS

RIVA-RANITIDINE (Ranitidine hydrochloride) is contraindicated for patients known to have hypersensitivity to ranitidine or to any ingredient in the formulation. For a complete listing, see DOSAGE FORMS, COMPOSITION AND PACKAGING.

WARNINGS AND PRECAUTIONS

Gastrointestinal

Gastric Ulcer

Treatment with a histamine H₂-antagonist may mask symptoms associated with carcinoma of the stomach and, therefore, may delay diagnosis of that condition. Accordingly, where gastric ulcer is suspected the possibility of malignancy should be excluded before therapy with ranitidine (ranitidine hydrochloride) is instituted.

Concomitant NSAID Use

Regular supervision of patients who are taking non-steroidal anti-inflammatory drugs concomitantly with ranitidine is recommended especially in the elderly and in those with a history of peptic ulcer. Baseline endoscopy and histological evaluation is necessary to rule out gastric carcinoma.

Endocrine and Metabolism

Cyanocobalamin (Vitamin B12) Deficiency

The prolonged use of H₂-receptor antagonists, may impair the absorption of protein-bound Vitamin B₁₂ and may contribute to the development of cyanocobalamin (vitamin B₁₂) deficiency.

Use in Patients with a History of Acute Porphyria

Rare clinical reports suggest that ranitidine may precipitate acute porphyric attacks. Therefore, ranitidine should be avoided in patients with a history of acute porphyria.

Renal

Use in Impaired Renal Function

Ranitidine is excreted via the kidneys and, in the presence of renal impairment, plasma levels of ranitidine are increased and elimination prolonged. Accordingly, it is recommended in such patients, to decrease the dosage of ranitidine by one half. Accumulation of ranitidine with resulting elevated plasma concentrations will occur in patients with renal impairment (creatinine clearance less than 50 mL/min); a recommended daily dose of oral ranitidine in such patients should be 150 mg.

Sexual Function/Reproduction

Fertility

There are no data on the effects of ranitidine on human fertility. There were no effects on male and female fertility in animal studies (see TOXICOLOGY).

Special Populations

Pregnant Women:

The safety of ranitidine in the treatment of conditions where a controlled reduction of gastric secretion is required during pregnancy has not been established. Reproduction studies performed in rats and rabbits have revealed no evidence of ranitidine induced impaired fertility or harm to the foetus. Ranitidine crosses the placenta. Nevertheless, if the administration of ranitidine is considered to be necessary, its use requires that the potential benefits be weighed against possible hazards to the patient and to the fetus.

Nursing Women:

Ranitidine is secreted in breast milk in lactating mothers but the clinical significance of this has not been fully evaluated. Like other drugs, ranitidine should only be used during nursing if considered essential.

Geriatrics (> 65 years of age):

Since malignancy is more common in the elderly, particular consideration must be given to this before therapy with ranitidine is instituted. Elderly patients receiving non-steroidal anti-inflammatory drugs concomitantly with ranitidine should be closely supervised.

As with all medication in the elderly, when prescribing ranitidine, consideration should be given to the patient's concurrent drug therapy. Sporadic cases of drug interaction have been reported in elderly patients involving both hypoglycaemic drugs and theophylline. The significance of these reports cannot be determined at present, as controlled clinical trials with theophylline and ranitidine have not shown interaction. Elderly patients may be at increased risk for confusional states and depression.

In patients such as the elderly, persons with chronic lung disease, diabetes or the immunocompromised, there may be an increased risk of developing community acquired pneumonia. A large epidemiological study showed an increased risk of developing community acquired pneumonia in current users of H₂ receptor antagonists versus those who had stopped treatment, with an observed adjusted relative risk increase of 1.63 (95% CI, 1.07-2.48)

Pediatrics (<18 years of age):

Experience with ranitidine products in children is limited. It has, however, been used successfully in children aged 8 to 18 years in oral doses up to 150 mg twice daily.

ADVERSE REACTIONS

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

The following adverse reactions have been reported as events in clinical trials or in the routine management of patients treated with ranitidine hydrochloride. A cause and effect relationship to ranitidine is not always established.

Central Nervous System:

Headache, sometimes severe; malaise; dizziness; somnolence; insomnia; vertigo; and reversible blurred vision suggestive of a change in accommodation. Isolated cases of reversible mental confusion, agitation, depression, hallucinations have been reported, predominantly in severely ill elderly patients. In addition, reversible involuntary movement disorders have been reported rarely.

Cardiovascular:

Isolated reports of tachycardia, bradycardia, premature ventricular beats, AV block have been noted. Asystole has been reported in very few individuals with and without predisposing conditions following IV administration and has not been reported following oral administration of ranitidine (see WARNINGS AND PRECAUTIONS, DOSAGE AND ADMINISTRATION).

Gastrointestinal:

Constipation, diarrhea, nausea/vomiting and abdominal discomfort/pain.

Hepatic:

In normal volunteers, transient and reversible SGPT and SGOT values were increased to at least twice the pretreatment levels in 6 of 12 subjects receiving ranitidine 100 mg q.i.d intravenously for seven days, and in 4 of 24 subjects receiving 50 mg q.i.d intravenously for five days. Therefore, it may be prudent to monitor SGOT and SGPT in patients receiving intravenous treatment for five days or longer and in those with pre-existing liver diseases. With oral administration, there have been occasional reports of hepatitis, hepatocellular or hepatocanalicular or mixed, with or without jaundice. In such circumstances, ranitidine should be discontinued immediately. These are usually reversible, but in exceedingly rare circumstances, death has occurred.

Renal:

Very rare cases of acute interstitial nephritis have been reported.

Musculoskeletal:

Rare reports of arthralgia and myalgia.

Haematologic

Blood count changes (leukopenia, thrombocytopenia) have occurred in a few patients. These are usually reversible. Rare cases of agranulocytosis or pancytopenia, sometimes with marrow hypoplasia or aplasia have been reported.

Endocrine:

No clinically significant interference with endocrine or gonadal function has been reported. There have been a few reports of breast symptoms and breast conditions (such as gynecomastia and galactorrhea).

Dermatologic:

Rash, including cases suggestive of mild erythema multiforme. Rare cases of vasculitis and alopecia have been reported.

Other:

Rare cases of hypersensitivity reactions (including chest pain, bronchospasm, fever, rash, eosinophilia, anaphylaxis, urticaria, angioneurotic edema, hypotension) and small increases in serum creatinine have occasionally occurred after a single dose. Acute pancreatitis and reversible impotence has been reported rarely.

DRUG INTERACTIONS

Overview

Ranitidine has the potential to affect the absorption, metabolism or renal excretion of other drugs. The altered pharmacokinetics may necessitate dosage adjustment of the affected drug or discontinuation of treatment.

Drug-Drug Interactions

Interactions occur by several mechanisms including:

1) Inhibition of cytochrome P450-linked mixed function oxygenase system:

Ranitidine at usual therapeutic doses does not potentiate the actions of drugs which are inactivated by this enzyme system such as diazepam, lidocaine, phenytoin, propranolol and theophylline. There have been reports of altered prothrombin time with coumarin anticoagulants (e.g., warfarin). Due to the narrow therapeutic index, close monitoring of increased or decreased prothrombin time is recommended during concurrent treatment with ranitidine.

2) Competition for renal tubular secretion:

Since ranitidine is partially eliminated by the cationic system, it may affect the clearance of other drugs eliminated by this route. High doses of ranitidine (e.g., such as those used in the treatment of Zollinger-Ellison syndrome) may reduce the excretion of procainamide and N-acetylprocainamide resulting in increased plasma levels of these drugs.

3) Alteration of gastric pH:

The bioavailability of certain drugs may be affected. This can result in either an increase in absorption (e.g., triazolam, midazolam) or a decrease in absorption (e.g. ketoconazole, atazanavir, delavirdine, gefitinib).

Hypoglycaemic drugs and theophylline: Sporadic cases of drug interactions have been reported in elderly patients involving both hypoglycaemic drugs and theophylline. The significance of these reports cannot be determined at present, as controlled clinical trials with theophylline and ranitidine have not shown interaction.

Sucralfate: If high doses (two grams) of sucralfate are co-administered with ranitidine, the absorption of ranitidine may be reduced. This effect is not seen if sucralfate is taken at least two hours after ranitidine administration.

Drug-Food Interactions

Interactions with food have not been established.

Drug-Herb Interactions

Interactions with herbal products have not been established.

Drug-Laboratory Test Interactions

There are no known interactions between ranitidine and laboratory tests.

DOSAGE AND ADMINISTRATION

Recommended Dose and Dosage Adjustment

Duodenal ulcer or benign gastric ulcer:

300 mg once daily at bedtime or 150 mg twice daily taken in the morning and before retiring. It is not necessary to time the dose in relation to meals. In most cases of duodenal ulcer and benign gastric ulcer, healing will occur in four weeks. In the small number of patients whose ulcers may not have fully healed, these are likely to respond to a further four week course of therapy. In the treatment of duodenal ulcers, 300 mg twice daily for 4 weeks may be of benefit when more rapid healing is desired.

Maintenance therapy:

Duodenal ulcers, benign gastric ulcers:

Patients who have responded to short-term therapy, particularly those with a history of recurrent ulcer, may benefit from chronic maintenance therapy at a reduced oral dosage of 150 mg once daily at bedtime.

In the management of duodenal ulcers, smoking is associated with a higher rate of ulcer relapse (up to 9.2 times higher in one trial), and such patients should be advised to stop smoking. In those patients who fail to comply with such advice, 300 mg nightly provides additional therapeutic benefit over the 150 mg once daily dosage regimen.

Reflux esophagitis:

Acute treatment:

300 mg once daily at bedtime, or alternatively 150 mg twice daily, taken in the morning and before retiring for up to eight weeks. In patients with moderate to severe esophagitis, the dosage of ranitidine may be increased to 150 mg four times daily for up to 12 weeks.

Long-term Management:

For the long-term management of reflux esophagitis, the recommended adult oral dose is 150 mg twice daily.

Post-operative peptic ulcer:

150 mg twice daily, taken in the morning and before retiring.

Pathological hypersecretory conditions (Zollinger-Ellison Syndrome):

150 mg three times daily may be administered initially. In some patients, it may be necessary to administer ranitidine 150 mg doses more frequently. Doses should be adjusted to individual patient needs. Doses up to six grams per day have been well tolerated.

Treatment of NSAID-induced lesions (both ulcers and erosions) and their gastrointestinal symptoms and prevention of their recurrence:

In ulcers following non-steroidal anti-inflammatory drug therapy or associated with continued non-steroidal anti-inflammatory drugs, 150 mg twice daily for 8-12 weeks may be necessary. For the prevention of non-steroidal anti-inflammatory drug associated ulcer recurrence, 150 mg twice daily may be given concomitantly with non-steroidal anti-inflammatory drug therapy.

Prophylaxis of acid aspiration syndrome (AAS):

150 mg the evening prior to anaesthesia induction is recommended, however, 150 mg two hours before anaesthesia induction is also effective. For the prevention of AAS in pre-partum patients who elect for anaesthesia, 150 mg every six hours may be employed, but if general anaesthesia is warranted, a non-Particulate oral antacid (e.g., sodium citrate) could supplement ranitidine therapy. In an emergency situation, the use of alkalis, antacids, and meticulous anaesthetic technique is still necessary as ranitidine does not affect the pH and volume of the existing gastric content.

Prophylaxis of hemorrhage from stress ulceration in seriously ill patients or prophylaxis of recurrent hemorrhage in patients bleeding from peptic ulceration who are currently managed by intravenous ranitidine:

An oral dose of 150 mg twice daily may be substituted for the injection once oral feeding commences.

Dosage for the Elderly:

For all conditions listed above, the drug dosage for the elderly who are seriously ill should start at

the lowest recommended dose and be adjusted as necessary with close supervision.

Patients over 50 years of age (see ACTIONS AND CLINICAL PHARMACOLOGY, Patients over 50 years of age).

OVERDOSAGE

There is no experience to date with deliberate overdosage. The usual measures to remove unabsorbed drug from the gastrointestinal tract (including activated charcoal or syrup of ipecac), clinical monitoring and supportive therapy should be employed.

For management of a suspected drug overdose, contact your regional Poison Control Centre.

ACTION AND CLINICAL PHARMACOLOGY

Ranitidine is an antagonist of histamine at gastric H₂-receptor sites. Thus, ranitidine inhibits both basal gastric secretion and gastric acid secretion induced by histamine, pentagastrin and other secretagogues. On a weight basis ranitidine is between 4 and 9 times more potent than cimetidine. Inhibition of gastric acid secretion has been observed following intravenous, intraduodenal and oral administration of ranitidine. This response is dose-related, a maximum response being achieved at an oral dose of 300 mg/day.

Pepsin secretion is also inhibited but secretion of gastric mucus is not affected. Ranitidine does not alter the secretion of bicarbonate or enzymes from the pancreas in response to secretin and pancreozymin.

Ranitidine is rapidly absorbed after oral administration of 150 mg ranitidine, peak plasma concentrations (300 to 550 ng/mL) occurred after 1 to 3 hours. Two distinct peaks or a plateau in the absorption phase result from reabsorption of drug excreted into the intestine. These plasma concentrations are not significantly influenced by the presence of food in the stomach at the time of the oral administration nor by regular doses of antacids.

Bioavailability of oral ranitidine is approximately 50% to 60%. Serum protein binding of ranitidine in man is in the range 10 to 19%. The elimination half-life is approximately 2 to 3 hours. The principal route of excretion is the urine (40% recovery of free and metabolized drug in 24 hours).

There is a significant linear correlation between the dose administered and the inhibitory effect upon gastric acid secretion for oral doses up to 300 mg. A plasma ranitidine concentration of 50 ng/mL has an inhibitory effect upon stimulated gastric acid secretion of approximately 50%. Estimates of the IC₅₀ range from 36 to 94 ng/mL. Following the administration of 150 mg ranitidine orally, plasma concentrations in excess of this lasted for more than 8 hours and after 12 hours, the plasma concentrations were sufficiently high to have a significant inhibitory effect upon gastric secretion. In patients with duodenal ulcer, 150 mg oral ranitidine every 12 hours significantly reduced mean 24-hour hydrogen ion activity by 69% and nocturnal gastric acid output by 90%. Furthermore, 300 mg oral ranitidine at night is as effective in reducing 24-hour intragastric acidity as 150 mg ranitidine given orally twice daily.

Following administration of 50 mg ranitidine injection intramuscularly, plasma concentrations in excess of 100 ng/mL were achieved within 5 minutes and remained above this level for 4 to 6 hours.

Intravenous infusion (rate: 0.125 mg/kg/hour) produced a rise of intragastric pH between 5.6 and 7.0 after 2 hours and maintained this level over the 24 hour period when administered to seriously ill patients. The volume of gastric secretion was reduced by more than 55%. Doubling the infusion rate to 0.25 mg/kg/hour produced no further increase in gastric acid inhibition.

A single 50 mg intravenous bolus dose of ranitidine injection produced significant acid inhibition 8 to 9 hours after administration. When 13 seriously ill patients with 2 or more risk factors (shock, sepsis, respiratory failure, jaundice, renal insufficiency or peritonitis) were treated with a 50 mg intravenous bolus dose of ranitidine injection followed by a continuous infusion of 0.2 mg/kg/hour, the number of 'at risk' days (gastric pH less than 3.5 at 3 consecutive four-hour aliquots) was approximately half that of placebo treated patients.

Tablets:

In respect of both 24-hour acidity and nocturnal acid output, an oral dose of ranitidine 150 mg twice daily was superior to cimetidine 200 mg three times daily and 400 mg at night ($p<0.001$ and $p<0.05$, respectively).

Treatment of volunteers with an oral dose of ranitidine 150 mg twice daily for 7 days did not cause bacterial overgrowth in the stomach.

Volunteers treated with an oral dose of ranitidine have reported no significant gastrointestinal or central nervous system side effects; moreover pulse rate, blood pressure, electrocardiogram and electroencephalogram were not significantly affected in man following ranitidine administration.

In healthy human volunteers and patients, ranitidine, when administered orally did not influence plasma levels of the following hormones: cortisol, testosterone, oestrogens, growth hormone, follicle-stimulating hormone, luteinizing hormone, thyroid-stimulating hormone, aldosterone or gastrin; although like cimetidine, ranitidine reduced vasopressin output. Treatment for up to 6 weeks with ranitidine 150 mg twice daily by mouth did not affect the human hypothalamic-pituitary-testicular-ovarian or -adrenal axes.

Patients over 50 years of age:

In patients over 50 years of age, half-life is prolonged (3 to 4 hours) and clearance is reduced, consistent with the age-related decline of renal function. However, systemic exposure and accumulation are 50% higher. This difference exceeds the effect of declining renal function, and indicates increased bioavailability in older patients.

STORAGE AND STABILITY

Store between 15°C and 30°C. Protect from light, excessive heat and moisture.

DOSAGE FORMS, COMPOSITION AND PACKAGING

150 mg Tablets:

Each white, round, biconvex, coated tablet, debossed with "RNT" logo on one side and "150" on the other side contains ranitidine hydrochloride equivalent to 150 mg ranitidine and the following non-medicinal ingredients: Colloidal Silicon Dioxide, Croscarmellose Sodium, Lecithin, Magnesium Stearate, Microcrystalline Cellulose, Polyvinyl Alcohol Part Hydrolyzed, Talc, Titanium Dioxide and Xanthan Gum. They are available in bottles of 60 and 250 tablets.

300 mg Tablets:

Each white, capsule-shaped, biconvex, coated tablet, debossed with "RNT" logo on one side and "300" on the other side contains ranitidine hydrochloride equivalent to 300 mg ranitidine and the following non-medicinal ingredients: Colloidal Silicon Dioxide, Croscarmellose Sodium, Lecithin, Magnesium Stearate, Microcrystalline Cellulose, Polyvinyl Alcohol Part Hydrolyzed, Talc, Titanium Dioxide and Xanthan Gum. They are available in bottles of 30 and 100 tablets.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION


Drug Substance

Proper Name: ranitidine hydrochloride

Chemical Name: (N-{2-[({5-[{(dimethylamino)-methyl]-2-furanyl}-methyl)thio]ethyl}-N'-methyl-2-nitro-1, 1-ethenediamine, hydrochloride

Molecular formula and molecular mass: C₁₃H₂₂N₄O₃S·HCl, 350.87g/mol (as hydrochloride salt)

Structural formula:



Physicochemical properties:

Ranitidine hydrochloride is a white to pale yellow granular substance. At room temperature, ranitidine hydrochloride is soluble in water, methanol, ethanol and chloroform (decreasing order).

pH and pKa values: pH (in 1% w/v aqueous solution): 5.40; pKa: 3.8

CLINICAL TRIALS

Comparative Bioavailability Studies

A single center, randomized, single-dose, blinded, 2-period, 2-sequence crossover comparative bioavailability study was performed under fasting conditions on Ranitidine Hydrochloride tablets using Laboratoire Riva Inc. RIVA-RANITIDINE 300 mg tablets (Lot # P-1417) versus the reference product, ZANTAC® 300 mg Tablets (Lot # 4A002), by GlaxoSmithKline Inc. Canada. Twenty-two healthy male volunteers took part in the study. The pharmacokinetic data calculated for the RIVA-RANITIDINE 300 mg and ZANTAC® 300 mg tablets formulation are tabulated below:

SUMMARY TABLE OF THE COMPARATIVE BIOAVAILABILITY DATA

Ranitidine (1 x 300 mg tablet) From measured data uncorrected for potency Geometric Mean Arithmetic Mean (CV %)				
Parameter	RIVA-RANITIDINE	ZANTAC®	% Ratio of Geometric Means	Confidence Interval (90%)
AUC _T (ng·h/mL)	4656.9 4801.8 (25.2)	4636.2 4775.4 (23.1)	100.45	94.03 – 107.31
AUC _I (ng·h/mL)	4781.1 4929.1 (25.0)	4759.3 4901.2 (23.0)	100.46	94.15 – 107.19
C _{max} (ng/mL)	987.2 1032.4 (30.5)	915.9 961.8 (32.4)	107.79	98.86 – 117.52
T _{max} [§] (h)	2.84 (1.33 – 4.00)	3.00 (1.67 – 4.00)		
T _½ [€] (h)	2.79 (11.7)	2.85 (16.4)		

[†]ZANTAC® was manufactured by GlaxoSmithKline Inc., Canada, and was purchased in Canada.

[§]Expressed as the median (range)

[€]Expressed as the arithmetic mean (CV %)

Study Results

In 6 clinical trials examining the healing of duodenal ulcers in 1,500 patients, a dose of 300 mg daily for 4 weeks was found to have an 83% healing rate; however, increasing the dose to 300 mg twice daily gave significantly better results (92% healed at 4 weeks; $p<0.001$).

DETAILED PHARMACOLOGY

Animal Pharmacology

Ranitidine is a potent competitive reversible, selective antagonist of histamine at H₂-receptors *in vitro* and *in vivo*. Thus, ranitidine antagonised the actions of histamine at H₂-receptors in the rat isolated uterus and in the guinea pig isolated atrium. Ranitidine is not an anticholinergic agent. On a molar basis, ranitidine is 4 to 5 times more active than cimetidine with a pA₂ value of 7.2. In concentrations 1,000 times greater than those required to block H₂-receptors, it failed to block either H₁-receptors or muscarinic receptors in the guinea pig isolated ileum. The beta-adrenoceptor responses of the rat uterus and guinea pig atrium to isoprenaline were also unaffected by ranitidine.

Blockade of histamine H₂-receptors in the stomach *in vivo* is the pharmacological action of ranitidine with greatest immediate clinical relevance. Ranitidine inhibits gastric secretion induced by various secretagogues in both the rat and dog.

In the conscious dog with a Heidenhain pouch, ranitidine given orally or intravenously antagonised gastric acid secretion induced by histamine, pentagastrin and bethanechol. Ranitidine was 5 to 10 times more active than cimetidine. However, both ranitidine and cimetidine had similar time curves of action. Ranitidine also inhibited the gastric secretory response to food in the conscious fistulated dog.

Ranitidine inhibited acid secretion in the perfused stomach of the anaesthetised rat, and acetylsalicylic acid-induced gastric lesion formation in the conscious rat, both in the presence and absence of excess hydrochloric acid. Measurements of the ratio of mucosal blood flow to acid secretion show that the inhibitory action of ranitidine upon gastric acid secretion cannot be attributed to changes in blood flow.

There were no behavioural effects in the mouse and rat after oral administration of 800 mg/kg ranitidine. Cats and dogs dosed with ranitidine 80 mg/kg orally, exhibited no behavioural effects indicative of an action on the central nervous system, although at this high dose level in the dog there was an indication of peripheral vasodilation and skin irritation due to released histamine. Ranitidine, when coadministered with the following CNS modulating preparations; codeine, hexobarbitone, ethyl alcohol, chlordiazepoxide, chlorpromazine, imipramine, α -methyldopa, reserpine, apomorphine or pentylenetetrazol, did not alter the pharmacological effects of either preparation.

At a dose level 45 times the antisecretory ED₅₀, intravenous infusion of ranitidine had no effect on the heart rate, blood pressure or electrocardiogram of the anaesthetised dog. The respiratory system was unaffected by ranitidine after oral doses in the mouse, rat, rabbit, cat and dog and after intravenous doses in the dog.

In the conscious dog, ranitidine had no appreciable effect on blood pressure or heart rate when administered orally at 10 mg/kg. There were short-lived falls in diastolic blood pressure after an intravenous dose of 10 mg/kg, 370 times the antisecretory dose level. There was no evidence of arrhythmia or of any electrocardiographic abnormality.

Long-term toxicity studies have shown that ranitidine does not possess antiandrogenic activity nor does it displace dihydrotestosterone from the androgen binding sites.

Metoclopramide, atropine and acetylsalicylic acid in the rat produced no change in the antisecretory activity of ranitidine.

The effect of ranitidine on anti-inflammatory drugs was varied. There was no effect on the anti-inflammatory action of prednisolone, but the anti-inflammatory action of indomethacin was enhanced. Administration of ranitidine reduced the frequency of acetylsalicylic acid- and indomethacin-induced gastric erosions. The antinociceptive action of acetylsalicylic acid was reduced after ranitidine treatment.

Ranitidine, unlike cimetidine, does not inhibit the hepatic mixed function oxygenase system. Spectral interaction studies have shown that whilst cimetidine binds strongly to cytochrome P₄₅₀, ranitidine has only weak affinity for this enzyme. Cimetidine is known to impair the metabolism of pentobarbitone and warfarin. In doses of up to 166 mg/kg in the rat, ranitidine had no effect on the pentobarbitone sleeping time or the pharmacokinetics and pharmacodynamics of warfarin.

Metabolism, Distribution and Excretion

The metabolism of ranitidine hydrochloride has been studied in four species of laboratory animal (mouse, rat, rabbit and dog) using radio-labelled drug. The drug was rapidly absorbed after oral administration. In the mouse, rat and rabbit between 30% and 60% of the administered radioactivity was excreted in the urine, the remainder being recovered in the feces.

In the mouse 47% was excreted in the urine within 24 hours. In the rat, N-demethylation of ranitidine was the major route of metabolism. 30% of the administered dose was excreted in the urine as unchanged drug, up to 14% as desmethylranitidine, 3-6% as the N-oxide and 4% as the S-oxide. In rat bile the major radioactive components were ranitidine and an unidentified metabolite known as "Fast-Running Metabolite" (FRM) which is thought to be a charge transfer complex of ranitidine with bile pigments.

In the rabbit, sulphoxidation of ranitidine was the major route of metabolism, 18% of the administered dose being excreted in the urine as unmetabolised ranitidine, 8% as S-oxide, 2-4% as the N-oxide, and 2-4% as desmethylranitidine.

In the dog up to 70% of the administered dose was excreted in the first 24 hours. About 40% of the drug was excreted in the urine as unchanged ranitidine and up to 30% as the N-oxide, N-oxidation being the main route of metabolism of ranitidine in the dog. The N-oxide was also the major radioactive component present in dog bile together with small amounts of unchanged ranitidine and FRM.

In the rat, rabbit and dog, less than 10.1% of ranitidine in plasma is protein bound. Within one to seven days of administration of radio-labelled drug in the rat and dog over 99% of the radioactivity was cleared from the body. In common with many drugs, radioactivity persisted in the uveal tract of these two species, the half-life in the dog uveal tract being of the order of 6 months. Ranitidine and its S-oxide have greater affinity for melanin than the desmethyl metabolite; the N-oxide is bound only to a small extent.

The placental transfer of radioactive ranitidine and its metabolites has been studied in the pregnant rat and rabbit. Whole body autoradiography of rat and rabbit foetuses showed that small amounts of radioactivity were present in the uveal tract of the foetal eye in both species, in the gall bladder and intestine of the rabbit foetus and in the bladder of the rat foetus. Radioactivity was also detected in the salivary and mammary glands of the maternal rat and at very low concentration, in the milk.

Human Pharmacokinetics

Serum concentrations necessary to inhibit 50% of stimulated gastric acid secretion are estimated to be 36 to 94 ng/mL. Following a single oral dose of 150 mg, serum concentrations of ranitidine are in this range for up to 12 hours. There is a relationship between plasma concentrations of ranitidine and suppression of gastric acid production but wide interindividual variability exists.

Ranitidine is 50% absorbed after oral administration compared to an IV injection with mean peak levels of 440 to 545 ng/mL occurring two to three hours after a 150 mg dose. The elimination half-life is 2 to 3 hours.

The major route of elimination is renal. After IV administration of 150 mg ³H-ranitidine, 98% of the dose was recovered, including 5% in feces and 93% in urine, of which 70% was unchanged parent drug. After oral administration of 150 mg ³-H ranitidine, 96% of the dose was recovered, 26% in the feces and 70% in urine of which 35% was unchanged parent drug. Less than 3% of the dose is excreted in bile. Renal clearance is approximately 500 mL/min, which exceeds glomerular filtration indicating net renal tubular secretion.

Ranitidine is absorbed very rapidly after an intramuscular injection. Mean peak levels of 576 ng/mL occur within 15 minutes or less following a 50 mg intramuscular dose. Absorption from intramuscular sites is virtually complete, with a bioavailability of 90% to 100% compared with intravenous administration.

The principal route of excretion is the urine, with approximately 30% of the orally-administered dose collected in the urine as unchanged drug in 24 hours. Renal clearance is about 530 mL/min, indicating active tubular excretion, with a total clearance of 760 mL/min. The volume of distribution ranges from 96 to 142 L.

Studies in patients with hepatic dysfunction (compensated cirrhosis) indicate that there are minor, but clinically insignificant alterations in ranitidine half-life, distribution, clearance and bioavailability.

Serum protein binding averages 15%.

The gastric antisecretory activity of ranitidine metabolites has been examined. In man, both the principal metabolite in the urine, the N-oxide (4% of the dose) and the S-oxide (1%) possess weak H₂-receptor blocking activity but desmethylranitidine (1%) is only 4 times less potent than ranitidine in the rat and half as potent as ranitidine in the dog.

TOXICOLOGY

Toxicology, Impairment of Fertility, Carcinogenesis, and Mutagenesis

Ranitidine hydrochloride has been subjected to exhaustive toxicological testing which has demonstrated the lack of any specific target organ or any special risk associated with its clinical use.

Non-clinical data revealed no special hazard for humans based on conventional studies of safety pharmacology, repeated-dose toxicity, genotoxicity, carcinogenic potential and toxicity to reproduction and development.

Acute Toxicity Studies

In mice and rats, the intravenous LD₅₀ is of the order of 75 mg/kg, whereas orally, even doses of 1000 mg/kg are not lethal. In dogs, the oral minimum lethal dose is 450 mg/kg/day. High single doses of ranitidine (up to 80 mg/kg orally) show only minimal and reversible signs of toxicity, some of which are related to transitory histamine releases.

Long-Term Toxicity Studies

In the long-term toxicity and carcinogenicity studies, very high doses of ranitidine were given daily to mice (up to 2000 mg/kg/day) throughout their normal life-span, and to dogs (up to 450 mg/kg/day) for periods of up to one year.

These doses produced massive plasma ranitidine concentrations far in excess of those found in human patients receiving ranitidine at the recommended therapeutic dose. For example, in the dogs, peak plasma concentrations were in excess of 115 mcg/mL and in mice basal plasma levels were in the range of 4-9 mcg/mL. In man, after oral administration of 150 mg ranitidine, the mean peak plasma concentration (C_{max}) was between 360 and 650 ng/mL.

In the rat, doses as high as 2000 mg/kg/day were well tolerated, the only morphological change seen was the increased incidence of accumulations of foamy alveolar macrophages in the lungs. The accumulation of these cells is a natural phenomenon in aging rats and chronic administration of a wide variety of drugs has been known to contribute to this process. Therefore, it is unlikely that the pharmacologic concentrations of ranitidine administered to these rats contributed to this natural process.

In the six-week and six-month oral studies in the dog (100 mg/kg/day) loose faeces were occasionally detected, while in the six-month study loose stools were accompanied on eight occasions by mucus-like material and sometimes by blood, mostly from one dog. Loose faeces, salivation and vomiting were observed in the 54-week dog study.

In isolated cases, dogs passed red-stained faeces which occasionally tested positive for occult blood. When the dose level was increased from 100 mg/kg/day to 225 - 450 mg/kg/day, no further red-stained faeces were seen, suggesting that any relationship to ranitidine is unlikely. Post-mortem examination of the dogs revealed no ranitidine-induced changes in the alimentary tract.

One dog had marginally raised levels of plasma alanine aminotransferase and alkaline phosphatase during the six-week study. This same dog also showed some necrotic foci in the liver. Small lesions of focal necrosis and fibrosis were also seen in one piece of liver from one female dog treated with 100 mg/kg for six months. No other differences were detected by light and electron microscopic examination of the treated and control livers. Since the focal lesions were seen in only one dog and were restricted to one piece of liver, it suggests that they were not caused by ranitidine.

Muscular tremors, an inability to stand, and rapid respiration were seen on occasion in dogs treated with 225 mg/kg/day in the 54-week study. The prevalence of these observations was increased when the dose was increased to a toxic level of 450 mg/kg/day. One dog died: no specific pathological changes or reason for the death was discovered.

Changes in the colour or granularity of the tapetum lucidum of the eye were detected in three dogs receiving the highest dose of ranitidine (450 mg/kg/day) during the 54-week study. In one dog this change was considered to be related to treatment. The change, a pallor of the tapetum, was reversible. No changes were seen with light or electron microscopic examination of the eye. The changes in the tapetum are of no clinical significance in humans since (i) humans do not have a tapetum lucidum and (ii) the changes were only seen at toxic pharmacological concentrations of ranitidine.

The mean serum glutamic pyruvic transaminase values for dogs treated at 450 mg/kg/day were significantly greater, albeit marginally, than the control values. These enzyme increases were not accompanied by any histological changes.

Studies in which ranitidine was administered parenterally were performed. No sign of specific local irritation attributable to ranitidine was detected. In the rat, no biochemical or histopathological changes were observed at intravenous dose levels as high as 20 mg/kg. Specifically, no significant changes were found in the veins or subcutis. Mild lesions in some muscle samples were observed: usually, the cells were basophilic and smaller than normal; and the nuclei were swollen, more numerous, and sometimes had migrated to the centre of the cell.

In the rabbit, slight infiltration of the pannicular muscle by mononuclear cells was noted. This minor subcutaneous reaction was uncommon and showed no group related distribution. There was no apparent difference in irritance between ranitidine injection and placebo injection. In the rat, intravenous ranitidine at dose levels of 5.0 and 10.0 mg/kg daily for 15 days and 28 days produced no treatment related changes of biological importance in the haematopoietic system.

In Beagle dogs, intravenous ranitidine injection in doses up to 10 mg/kg/day for 28 and 42 days, produced no drug-related change in circulating erythrocytes or leukocytes and had no adverse effects on the haematopoietic system. No dose related changes were seen in electrocardiograms of Beagle dogs receiving up to 10 mg/kg ranitidine by intravenous injection. At dosage levels of up to

30 mg/kg, administered twice daily to Beagle dogs for 14 or 15 days, intravenous ranitidine injection produced no changes of biological significance in haematology, clinical chemistry or urinalysis.

No changes were observed in the eyes of dogs (specifically the tapetum lucidum) receiving ranitidine in doses up to 30 mg/kg twice daily for 15 days. At intravenous doses above 1.25 mg/kg, ranitidine injection produced immediate and transient reactions in the Beagle dog. The following reactions were typically produced by the administration of 1.25 mg/kg: bloodshot eyes, closing and watering of eyes, defaecation, diarrhoea, erythema, flatus, licking of lips, running nose, salivation, subdued behaviour, swallowing, tachycardia, and trembling. The range and severity of the effects was aggravated by increased dosage.

Reproduction Studies (Impairment of Fertility)

Reproduction studies were carried out in the rat and rabbit.

Rats were exposed to ranitidine before and during mating, throughout pregnancy, lactation and during the weaning period. No effects on the reproductive process were seen and there was no evidence of an anti-androgenic effect.

A total of 2,297 foetuses from rats treated with ranitidine were examined. There was no evidence that ranitidine is a rat teratogen. Cleft palates occurred in foetuses from both treatment groups, however, there were significantly more in the control rat population.

A total of 944 foetuses from rabbits treated with ranitidine were examined; no drug-related adverse events or abnormalities in the foetuses were observed.

Rabbits receiving a bolus intravenous injection of ranitidine (10 mg/kg) once daily on gestation days 7-16 exhibited a reduction in weight gain. Their foetuses weighed significantly less than foetuses of untreated controls. In addition, 12.4% of ranitidine-exposed foetuses had cleft palates. Reanalysis of this and a companion study performed to assess reproducibility demonstrated a lack of data reproducibility. Therefore, the effects observed in the first trial are aberrant, and should not form the basis for maternal or fetal toxicity.

In the subsequent study, no evidence of maternal or foetal toxicity was observed in rabbits dosed with 100 mg/kg ranitidine orally during days 2-29 of pregnancy. The peak plasma levels of ranitidine after a 100 mg/kg oral dose are similar to those obtained one minute after a 10 mg/kg dose administered intravenously (20-25 mcg/mL). Therefore, no teratogenic effects of ranitidine have been demonstrated at doses of 10 mg/kg (IV) and 100 mg/kg (Tablets) in rabbits.

Carcinogenicity Studies

There is no evidence that ranitidine is a carcinogen. Long term toxicity and carcinogenicity studies have involved the treatment of 600 mice and 636 rats at doses up to 2,000 mg/kg for two years and 129 weeks respectively and 42 dogs at doses up to 450 mg/kg/day for periods up to one year. These dose levels are far in excess of those to be used therapeutically in man. None of these animals had

any intestinal metaplasia. There was no evidence of a tumorigenic effect of ranitidine in any other tissue.

Mutagenesis

Ranitidine is not mutagenic at doses as great as 30 mg/plate in the Ames Assay utilizing *Salmonella typhimurium* (TA 1538, TA 98, TA 100 and TA 1537) or in doses of 9 mg/plate utilizing *Escherichia coli* (WP2 and WP2 uvrA) with or without activation.

Ranitidine at concentrations of 20-30 mg/plate had a weak direct mutagenic action in *S. typhimurium* TA 1535 and at 9 mg/plate in *E. coli* WP67. Ranitidine was not mutagenic at a concentration of 2 mg/mL in *E. coli* or *S. typhimurium* in the more sensitive Oral Solution microtitre fluctuation assay method. This weak direct mutagenic effect is of no clinical significance; the magnitudes of ranitidine concentration used in these assays are thousands of times greater than that attained therapeutically in human plasma.

The principal metabolites of ranitidine in man were not significantly mutagenic. This conclusion is supported by the following experiment. A test solution obtained by interacting ranitidine (10 mM) and sodium nitrite (40 mM) was mutagenic in *S. typhimurium* (TA 1535) but not in *S. typhimurium* (TA 1537) or in *E. coli* (WP67 or WP2 uvrA). This positive result is attributable to the presence of a nitrosonitrolic acid derivative AH 23729, which was mutagenic. When the sodium nitrite concentration was reduced to 15 mM or less, the solution was not mutagenic in any of the test microorganisms. The formation of AH 23729 requires concentrations of nitrous acid far in excess of those encountered in any probable physiological conditions. The other nitrosation products were not mutagenic in any of the microorganisms tested. There is no reason, therefore, for supposing that ranitidine is likely to be mutagenic in animals or man as a consequence of nitrosation in the stomach.

There is no evidence from long term toxicology, carcinogenicity and mutagenicity studies in animals to suggest that ranitidine is likely to have any deleterious effects in man when administered at therapeutic dose levels.

REFERENCES

1. Ashton MG, Holdsworth CD, Ryan FP, Moore M. Healing of gastric ulcers after one, two and three months of ranitidine. *Br. Med. J.* 1982; 284: 467-468.
2. Bell JA, Dallas FAA, Jenner WN, Martin LE. The metabolism of ranitidine in animals and man. [Abstract] *Biochem. Soc. Trans.* 1980; 8: 93.
3. Bories P, Michel H, Duclos B, Beraud JJ, Mirouse J. Use of ranitidine without mental confusion in patients with renal failure. [Letter] *Lancet* 1980; 2: 755.
4. Boyd EJ, Wilson JA, Wormsley KG. Review of ulcer treatment: role of ranitidine. *J. Clin. Gastroenterology* 1983; 5 Suppl 1:133-141.
5. Breen KJ, Bury RD, Desmond PV, et al. Effects of cimetidine and ranitidine on hepatic drug metabolism. *Clin. Pharmacol. Ther.* 1982; 31: 297-300.
6. Brogden RN, Carmine AA, et al. Ranitidine: A review of its pharmacology and therapeutic use in peptic ulcer disease and other allied diseases. *Drugs* 1982; 24: 267-303.
7. Critchlow JF. Comparative efficacy of parenteral histamine H₂ antagonists in acid suppression for the prevention of stress ulceration. *Am. J. Med.* 1987; 83: 23-28.
8. Damman HG, Muller P, Simon B. Parenteral ranitidine: onset and duration of action. *Br. J. Anaesth.* 1982; 54: 1235-1236.
9. Danilewitz M, Ou Tim L, Hirschowitz B. Ranitidine suppression of gastric hypersecretion resistant to cimetidine. *N. Engl. J. Med.* 1982; 306: 20-22.
10. Domschke W, Lux G, Domschke S: Furan H₂-antagonist ranitidine inhibits pentagastric-stimulated gastric secretion stronger than cimetidine. *Gastroenterology* 1980; 79: 1267-1271.
11. Durrant JM, Strunin L. Comparative trial of the effect of ranitidine and cimetidine on gastric secretion in fasting patients at induction of anaesthesia. *Can. Anaesth. Soc. J.* 1982; 29: 446-451.
12. Ehsanullah RSB, Page MC, Tildesley G, Wood JR. A placebo-controlled study of ranitidine in healing NSAID-associated gastric and duodenal ulcers. *Br. J. Rheumatol.* 1990; 29 (Suppl. 2): 9, A17.
13. Freston JW. H₂-receptor antagonists and duodenal ulcer recurrence: analysis of efficacy and commentary on safety, costs and patient selection. *Am. J. Gastroenterol.* 1987; 82: 1242-1249.
14. Gaginella TS, Bauman JH. Ranitidine hydrochloride. *Drug Intell. Clin. Pharm.* 1983; 17: 873-885.
15. Goudsouzian NG, Young ET. The efficacy of ranitidine in children. *Acta Anaesthesiologica Scand.* 1987; 31: 387-390.

16. Halparin L, Reudy J. Inhibition of pentagastrin-stimulated gastric acid secretion by ranitidine hydrochloride and cimetidine. *Curr. Ther. Res.* 1980; 28: 154-162.
17. Harris PW, Morison DH, Dunn GL, et al. Intramuscular cimetidine and ranitidine as prophylaxis against gastric aspiration syndrome - a randomized double blind study. *Can. Anaesth. Soc. J.* 1984; 31: 599 - 603.
18. Jensen RT, Collen JM et al. Cimetidine-induced impotence and breast changes in patients with gastric hypersecretory states. *N. Engl. J. Med.* 1983; 308:883.
19. Knodell RG, Holtzman JL, Crankshaw DL et al. Drug metabolism by rat and human hepatic microsomes in response to interaction with H₂ receptor antagonists. *Gastroenterology* 1982; 82: 1007.
20. Konturek SJ, Obtulowicz W, Kwiecien N, Sito E, Mikos K, Olesky J. Comparison of ranitidine and cimetidine in the inhibition of histamine, sham-feeding and meal-induced gastric secretion in duodenal ulcer patients. *Gut* 1980; 21: 181-186.
21. Lancaster-Smith MJ, Jaderberg MA, Jackson DA. Ranitidine in the treatment of NSAID-associated gastric and duodenal ulcers. *Gut* 1991; 32:252-255.
22. Lebert PA, Mahon WA, et al. Ranitidine kinetics and dynamics II. Intravenous dose studies and comparison with cimetidine. *Clin. Pharmacol. Ther.* 1981; 30: 545-550.
23. Leeder JS, Tesoro AM, Bertho-Gebara CE, MacLeod SM. Comparative bioavailability of ranitidine Tablets and suspension. *Canadian Journal of Hospital Pharmacy.* 1984; 37(3), 92-94, 106.
24. Maile CJD, Francis RN. Pre-operative ranitidine. *Anaesthesia* 1983; 38: 324-326.
25. Misiewicz JJ, Sewing K. (eds.). Proceedings of the First International Symposium on Ranitidine. *Scand. J. Gastroenterol.* 1981; 16 (Suppl. 69): 1-131.
26. Misiewicz JJ, Wormsley KG (eds.). The Clinical Use of Ranitidine. The Medicine Publishing Foundation Symposium Series 5, Pembroke House, Oxford, 1982.
27. Neils GF, van de Meene JGC. Comparative effect of cimetidine and ranitidine on prolactin secretion. *Postgrad. Med. J.* 1980; 56: 478-480.
28. Page M, Lacey L. Ranitidine syrup in the treatment of duodenal ulcer. *American Journal of Gastroenterology.* 1987; 82(9), 977.
29. Pasquali R, Corinaldesi R, Miglioli M, et al. Effect of prolonged administration of ranitidine on pituitary and thyroid hormones, and their response to specific hypothalamic-releasing factors. *Clin. Endocrinol.* 1981; 15: 457-462.

30. Peden NR, Robertson AJ, Boyd EJS, et al. Mitogen stimulation of peripheral blood lymphocytes of duodenal ulcer patients during treatment with cimetidine or ranitidine. Gut 1982; 23: 398-403.
31. Riley AJ, Salmon PR (eds.). Ranitidine. Excerpta Medica, Amsterdam, 1982.
32. Roberts CJC. Clinical Pharmacokinetics of Ranitidine. Clin. Pharmacokin. 1984; 9: 211-221.
33. Scarpignato C, Bertaccine G, Zimbara G, Vitulo F. Ranitidine delays gastric emptying of solids in man. Br. J. Clin. Pharmacol. 1982; 13: 252-253.
34. Wolfe MM. Considerations for selection of parenteral histamine (H_2)-receptor antagonists. Am. J. Med. 1987; 83: 82-88.
35. Yeomans ND, Hanson RG, Smallwood RA, Mihaly GW, Louis WJ. Effect of chronic ranitidine treatment on secretion of intrinsic factor. Br. Med. J. 1982; 285:264.
36. Product Monograph for ^{Pr}ZANTAC[®] GlaxoSmithKline Inc., Mississauga, Ontario, Canada; Date of Revision: January 15, 2015; Control No. 179097.
37. Single Center, Randomized, Single Dose, Fasting, Bioequivalence Study of pms-Ranitidine 300 mg Tablets Versus ZANTAC[®] 300 mg Tablets in normal healthy male volunteers.

IMPORTANT: PLEASE READ

PART III: CONSUMER INFORMATION

RIVA-RANITIDINE Ranitidine tablets, USP 150 mg & 300 mg

This leaflet is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about RIVA-RANITIDINE. Contact your doctor or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATION

What the medication is used for:

- to heal ulcers in the stomach, or the part that it empties into (the duodenum).
- to prevent stomach ulcers which may be caused by medicines called non-steroidal anti-inflammatory drugs (NSAIDs), often used to treat arthritis
- to prevent ulcers from bleeding
- to heal or stop problems caused by acid in the food pipe (esophagus) or too much acid in the stomach. This can cause pain or discomfort sometimes known as indigestion or heartburn
- to stop acid coming up from the stomach while under anesthetic during an operation

What it does:

RIVA-RANITIDINE belongs to a group of medicines called H₂-receptor blockers. It works by reducing the amount of acid in your stomach.

When it should not be used:

Don't take RIVA-RANITIDINE if you are allergic (hypersensitive) to ranitidine or any other ingredients of RIVA-RANITIDINE (see What the non-medicinal ingredients are).

What the medicinal ingredient is:

Ranitidine hydrochloride

What the non-medicinal ingredients are:

Colloidal Silicon Dioxide, Croscarmellose Sodium, Lecithin, Magnesium Stearate, Microcrystalline Cellulose, Polyvinyl Alcohol Part Hydrolyzed, Talc, Titanium Dioxide and Xanthan Gum

What dosage forms it comes in:

Tablets: 150 mg and 300 mg

WARNINGS AND PRECAUTIONS

BEFORE you use RIVA-RANITIDINE talk to your doctor or pharmacist if you:

- have stomach cancer
- have kidney disease, your doctor may lower your dose of RIVA-RANITIDINE
- have a rare condition called acute porphyria (a blood disease)
- have lung disease

- are diabetic
- have any problems with your immune system
- have had stomach ulcers before and you are taking Non-Steroidal Anti-Inflammatory (NSAID) medicines
- are pregnant, planning to become pregnant, breastfeeding or planning to breastfeed
- are taking any other medications including NSAIDs (see Interactions with this Medication).

Under rare circumstances supervised by the doctor, H₂-receptor antagonists such as RIVA-RANITIDINE might be used for long periods. Long term use of H₂-receptor antagonists may prevent normal absorption of vitamin B₁₂ from the diet and could lead to vitamin B₁₂ deficiency. Talk to your doctor.

INTERACTIONS WITH THIS MEDICATION

Tell your doctor or pharmacist if you're taking any other medicines, if you've taken any recently, or if you start taking new ones. This includes medicines bought without a prescription. Some medicines can affect how RIVA-RANITIDINE works, or make it more likely that you'll have side effects. RIVA-RANITIDINE can also affect how some other medicines work.

Drugs that may interact with RIVA-RANITIDINE include:

- procainamide or n-acetylprocainamide (used to treat heart problems)
- warfarin (used to thin the blood)
- triazolam (used to treat insomnia)
- midazolam (a sedative that may be given just before an operation)
- ketoconazole (used to treat fungal infections)
- atazanavir or delavirdine (used to treat HIV)
- gefitinib (used to treat lung cancer)
- Non-Steroidal Anti-Inflammatory (NSAID) medicines (used to treat pain and inflammation)
- sucralfate (used to treat ulcers). Your doctor may advise that you take high doses or oral sucralfate (e.g., 2 g) at least 2 hours after RIVA-RANITIDINE administration.

PROPER USE OF THIS MEDICATION

Usual Adult Dose:

Always take RIVA-RANITIDINE exactly as your doctor has told you to. Check with your doctor or pharmacist if you're not sure.

The usual dose is either:

- 150 mg in the morning and 150 mg in the evening or,
- 300 mg at bedtime

Your exact dose will depend on your particular stomach condition. Your doctor will tell you the dose you should take.

Swallow each tablet whole with some water.

Overdose:

IMPORTANT: PLEASE READ

If you take too much RIVA-RANITIDINE, contact your doctor or pharmacist for advice. If possible, show them the RIVA-RANITIDINE pack.

In case of drug overdose, contact a health care practitioner, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Side effects may include:

- allergic reactions
- skin rash
- inflammation of blood vessels (vasculitis)
- inflammation of the pancreas (pancreatitis)
- inflammation of the liver (hepatitis), sometimes with yellowing of the whites of the eyes or skin (jaundice)
- inflammation in the kidney (interstitial nephritis)
- slow, fast or irregular heartbeat
- diarrhea, constipation, nausea, vomiting, stomach pain
- feeling confused, depressed, or excited, or seeing or hearing things that are not really there (hallucinations), trouble sleeping (insomnia); feeling sleepy (somnolence)
- joint or muscle pain, malaise, uncontrolled movement
- headache, dizziness, blurred vision
- unusual hair loss or thinning (alopecia)
- unable to get or maintain an erection (impotence)
- unusual secretion of breast milk or breast enlargement in men

If you have any concerns about the side effects, tell your doctor, nurse or pharmacist.

Side effects that may show up in your blood tests:

- changes to liver function
- low levels of white blood cells
- decrease in number of blood platelets (cells that help blood to clot)
- decrease in number of all types of blood cells
- small increase in the level of creatinine (a waste product) in your blood

SERIOUS SIDE EFFECTS , HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom/effect	Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
	Only if severe	In all cases	
<u>Hypersensitivity</u> <u>Reaction</u> raised and itchy rash (hives), swelling, sometimes of the face or mouth (angioedema), chest pain, shortness of breath, unexplained fever, wheezing or difficulty in breathing, feeling faint, especially when standing up, collapse			✓
<u>Serious Skin Reactions</u> Skin rash, which may blister, and look like small targets (central dark spots surrounded by a paler area, with a dark ring around the edge)			✓
<u>Hepatitis</u> Yellowing of the skin or whites of the eyes, dark or tea colored urine, pale colored stools/ bowel movements, nausea/ vomiting, loss of appetite, pain, aching or tenderness on right side below the ribs			✓
<u>Cardiovascular</u> Slow, fast or irregular heartbeat			✓

This is not a complete list of side effects. If you have any unexpected effects while taking this RIVA-RANITIDINE, contact your doctor or pharmacist.

HOW TO STORE IT

IMPORTANT: PLEASE READ

Store RIVA-RANITIDINE between 15°C and 30°C. Protect from light, excessive heat and moisture. Keep out of reach and sight of children.

REPORTING SIDE EFFECTS

You can help improve the safe use of health products for Canadians by reporting serious and unexpected side effects to Health Canada. Your report may help to identify new side effects and change the product safety information.

3 ways to report:

- Online at MedEffect™ (www.healthcanada.gc.ca/medeffect);
- By calling 1-866-234-2345 (toll-free);
- By completing a Consumer Side Effect Reporting Form and sending it by:

- Fax to 1-866-678-6789 (toll-free), or
- Mail to: Canada Vigilance Program
Health Canada, Postal Locator
1908C
Ottawa, ON
K1A 0K9

Postage paid labels and the Consumer Side Effect Reporting Form are available at MedEffect™ (www.healthcanada.gc.ca/medeffect).

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

This document plus the full product monograph, prepared for health professionals can be obtained by contacting Laboratoire Riva Inc., at 1-800-363-7988.

This leaflet was prepared by
Laboratoire Riva Inc.
Blainville Québec
J7C 3V4

www.labriva.com

Last revised: February 3, 2017.